

BOSPHORUS MEDICAL JOURNAL

Boğaziçi Tıp Dergisi



ORIGINAL ARTICLES

Early Outcomes After Arthroscopic Rotator Cuff Repair Under General Anesthesia or Interscalene Block
Arif Çakar et al.

Assessment of Fall Risk
Elem Yorulmaz et al.

Performance of 'Turkish Triage System
Selman Faruk Yılmaz et al.

LLLT and Ultrasound in Shoulder Pain
Banu Aydeniz and Turan Uslu

Greater Occipital Nerve Block in Medication Overuse Headache
Ahmet Akpınar

CASE REPORT

◀ An Early-Stage Neuroborreliosis Case with Atypical Findings
Sema Yapıcı et al.

ISSN 2149-0287

Volume: 13 Number: 2 Year: 2026

BOSPHORUS MEDICAL JOURNAL

Boğaziçi Tıp Dergisi

EDITOR

Eren Gözke, M.D.

Department of Neurology, Fatih Sultan Mehmet Training and Research Hospital, İstanbul, Türkiye

ASSOCIATE EDITORS

Bekir Eray Kılınc, M.D.

Department of Orthopedics and Traumatology, Fatih Sultan Mehmet Training and Research Hospital, İstanbul, Türkiye

Esmâ Esin Derin Çiçek, M.D.

Department of Radiology, Fatih Sultan Mehmet Training and Research Hospital, İstanbul, Türkiye

Feyza Ünlü Özkan, M.D.

Department of Physical Medicine and Rehabilitation, Fatih Sultan Mehmet Training and Research Hospital, İstanbul, Türkiye

Neşe Keser, M.D.

Department of Brain and Neurosurgery, Fatih Sultan Mehmet Training and Research Hospital, İstanbul, Türkiye

Pelin Doğan Ak, M.D.

Department of Neurology, Fatih Sultan Mehmet Training and Research Hospital, İstanbul, Türkiye

Serkan Akan, M.D.

Department of Urology, Fatih Sultan Mehmet Training and Research Hospital, İstanbul, Türkiye

FOREIGN LANGUAGE EDITORS

Burcu Seher Anıl, M.D.

Department of Radiology, University of Health Sciences, Fatih Sultan Mehmet Training and Research Hospital, İstanbul, Türkiye

Ertunç Mega, M.D., Ph.D.

Department of Obstetrics and Gynecology, University of Health Sciences, Fatih Sultan Mehmet Training and Research Hospital, İstanbul, Türkiye

CONTACT

Eren Gözke, M.D.

Address: Department of Neurology, University of Health Sciences, Fatih Sultan Mehmet Training and Research Hospital, İstanbul

Phone: +90 216 578 30 00

Web: www.bosphorusmedj.com

E-mail: egozke@hotmail.com

OWNER

Barış Yılmaz, M.D.

On Behalf of University of Health Sciences, Fatih Sultan Mehmet Training and Research Hospital, İstanbul

PUBLISHER

Kare Publishing

Address: Göztepe Mah. Fahrettin Kerim Gökay Cad. No: 200 Da: 2, Göztepe, Kadıköy, İstanbul.

Phone: +90 216 550 61 11 Web: kareyayincilik.com E-mail: kare@kareyayincilik.com

Publications Coordinator: Suğra Türksavaş

Graphic Design: Zeynep Koyun

Bosphorus Medical Journal is indexed in the Open Ukrainian Citation Index, Scilit, EBSCO, CINAHL, Türkiye Citation Index, DOAJ, ProQuest, GALE, Research4life, Hinari and Idealonline.

© Copyright 2026 by Bosphorus Medical Journal - Available online at www.bosphorusmedj.com

BOSPHORUS MEDICAL JOURNAL

Boğaziçi Tıp Dergisi

AIMS & SCOPE

DESCRIPTION

Boğaziçi Tıp Dergisi – Bosphorus Medical Journal is a peer-reviewed journal published three times a year. The journal aims rapid publication of papers in all fields of the medical sciences on a regular basis. The journal publishes editorials, original papers of experimental and clinical research, reviews, case reports, letters to the editor and articles on the current issues of medicine.

AIM AND SCOPE

Bosphorus Medical Journal is the scientific open access publication of University of Health Sciences, Fatih Sultan Mehmet Training and Research Hospital (the content may be accessed free of charge through the web site www.bosphorusmedj.com). Three issues are released every year in April, August and December. The language of publication is English and Turkish.

BASIC PUBLICATION RULES

Authors are responsible for the accuracy of data and the journal is in compliance with the uniform requirements for manuscripts submitted to biomedical journals published by the ICMJE. The editorial and publication processes of the journal are conducted in accordance with the guidelines of the World Association of Medical Editors (WAME), the Council of Science Editors (CSE), the European Association of Science Editors (EASE), and the Committee on Publication Ethics (COPE) as well.

Statements or opinions expressed in the manuscripts published in Bosphorus Medical Journal reflect the views of the author(s) and not the opinions of the editors, the editorial board and the publisher; the editors, the editorial board and the publisher disclaim any responsibility or liability for such materials.

ABSTRACTING AND INDEXING

Bosphorus Medical Journal is indexed in the Open Ukrainian Citation Index, Scilit, EBSCO (2022), CINAHL (2022), Türkiye Citation Index (2020), DOAJ (2021), ProQuest (2021), GALE (2023), Research4life (2021), Hinari (2021) and Idealonline.

ABBREVIATION

Bosphorus Med J

PUBLICATION FEE

There is no fee for article submission, article processing or publication. Manuscripts can be reached from the web page of journal without any fees. No additional fee is required from the authors for accepted manuscripts.

Boğaziçi Tıp Dergisi / Bosphorus Medical Journal is a double-blind peer-reviewed, and open-access journal. The journal aims rapid publication of papers in all fields of the medical sciences on a regular basis. Boğaziçi Tıp Dergisi / Bosphorus Medical Journal publishes editorials, original papers of experimental and clinical research, reviews, case reports, letters to the editor and articles on the current issues of medicine.

Four issues are released every year in April, August and December. The language of publication is English. Free full-text articles are available at www.bosphorusmedj.com.

EDITORIAL POLICY

The editorial and publication processes of the journal are shaped in accordance with the guidelines of the International Council of Medical Journal Editors (ICMJE), the World Association of Medical Editors (WAME), the Council of Science Editors (CSE), the Committee on Publication Ethics (COPE), the European Association of Science Editors (EASE), and the National Information Standards Organization (NISO). The journal conforms to the Principles of Transparency and Best Practice in Scholarly Publishing (doaj.org/bestpractice).

Submission is a representation that neither the manuscript nor its data have been previously published (except abstracts and press reports published in connection with scientific meetings) or are currently under consideration for publication.

Materials copied from other sources must be accompanied by a written statement from both author and publisher giving permission to **Boğaziçi Tıp Dergisi / Bosphorus Medical Journal** for reproduction. It is the author's responsibility to ensure that permissions are obtained. The differences between the papers should be explained in detail.

Ethics committee approval of research protocols in accordance with international agreements (World Medical Association Declaration of Helsinki "Ethical Principles for Medical Research Involving Human Subjects," amended in October 2013, www.wma.net) is required for experimental, clinical, and drug studies, and for some case reports. If required, ethics committee reports or an equivalent official document will be requested from the authors.

For manuscripts concerning experimental research on humans, a statement should be included indicating that the written, informed consent of patients and volunteers was obtained following a detailed explanation of the procedures that they might undergo. For studies carried out on animals, the measures taken to prevent pain and suffering of the animals should be stated clearly.

Information on patient consent, the name of the ethics committee, and the ethics committee approval number should also be stated in the Materials and Methods section of the manuscript. It is the authors' responsibility to carefully protect the patients' anonymity. If there is a chance that a patient may be identified from a photograph or other image, or any accompanying text, releases signed by the patient or their legal representative should be enclosed. Identifying images should also be anonymized.

All submissions are screened with similarity detection software (iThenticate).

In the event of alleged or suspected research misconduct, e.g., plagiarism, citation manipulation, or data falsification/fabrication, the Editorial Board will follow up and act in accordance with COPE guidelines.

AUTHORSHIP CRITERIA

Boğaziçi Tıp Dergisi / Bosphorus Medical Journal follows the authorship criteria developed by the International Committee of Medical Journal Editors. All persons designated as authors should qualify for authorship, and all those who qualify should be listed. Each author should have participated sufficiently in the work to take public responsibility for appropriate portions of the content. One or more authors should take responsibility for the integrity of the work as a whole, from inception to published article.

Authorship credit should be based only on; (1) substantial contributions to conception and design, or acquisition of data, or analysis and interpretation of data, (2) drafting the article or revising it critically for important intellectual content, and (3) final approval of the version to be published. Conditions 1, 2, and 3 must all be met. Acquisition of funding, the collection of data, or general supervision of the

research group, by themselves, do not justify authorship. All others who contributed to the work who are not listed as authors should be named in the acknowledgments, and their contribution should be described. Authorship of multicenter trials is attributed to a group. All members of the group who are named as authors should fully meet the above criteria for authorship. Group members who do not meet these criteria should be listed, with their permission, in the acknowledgments. Financial and material support should also be acknowledged.

All statements and opinions expressed in the manuscripts published in Boğaziçi Tıp Dergisi / Bosphorus Medical Journal reflect the views of the author(s). All liability for the advertisements rests with the appropriate organization(s). The Editor-in-Chief, and Kare Publishing do not accept any responsibility for claims made in articles or advertisements.

The authors transfer all copyright to the manuscript to the Boğaziçi Tıp Dergisi / Bosphorus Medical Journal according to the framework of national and international regulations at the start of the evaluation process. A Copyright Transfer Form signed by the corresponding author must be submitted to the journal with the manuscript. A separate form should be submitted for each manuscript. Manuscripts submitted without a Copyright Transfer Form will not be accepted. After acceptance of the manuscript, all of the authors must complete and sign a Copyright Transfer Form. The authors must affirm that they will not submit the work to another journal, publish it in the original or another language, or allow a third party to use the manuscript without the written permission of the Boğaziçi Tıp Dergisi / Bosphorus Medical Journal. In the event that the manuscript is rejected, all copyrights transfer back to the authors. Manuscripts that are declined will not be returned, with the exception of artwork.

All content is the authors' responsibility. All financial liability and legal responsibility associated with the copyright of submitted tables, figures, and other visual materials protected by national and international laws rest with the authors. The authors take responsibility for any legal proceedings issued against the journal.

To clarify scientific contributions and responsibilities and any conflict of interest issues relevant to the manuscript, all parts of the Authors' Contribution Form must be completed by the corresponding author and the ICMJE Uniform Disclosure Form for Potential Conflicts of Interest must be completed online by all authors. Both forms should be included with the manuscript at the time of original submission.

Author names will be published as they are listed on the Copyright Transfer Form. Requests for changes in affiliation or the order of names at a later date cannot be granted to protect all parties involved.

REVIEW PROCESS

The journal aims at rapid publication of the submitted manuscripts. **Boğaziçi Tıp Dergisi / Bosphorus Medical Journal** promotes expert refereeing by peers as a tried and true method for the maintenance of standards of excellency in the scientific community. All papers are double blind peer-reviewed to determine the originality, validity and importance of content and conclusions. Manuscripts will be reviewed by two external reviewers for content, originality, importance to the field, appropriateness of statistical analysis, and derivation of conclusions. In the case of discrepancies between peer reviewers, the manuscript would be sent to a third reviewer. All reviewers remain anonymous. Final decision regarding the acceptance of the submitted articles belong to editor-in-chief.

All reviews will be completed within one month of submission to the journal. Authors will be sent reviewers comments that are judged to be useful to them. In principle, the instructions, objections and requests made by the reviewers should be strictly followed. With revised form of manuscript, the authors should state clearly and precisely every step taken in accordance with the reviewers' requests. The description should be listed on a numbered basis, in the order of reviewers' comments. Altered paragraphs in the new version of the manuscript should be specified using page and paragraph numbers. Paragraph on top of a page is considered no 1, even if it does not begin on that page.

All manuscript submissions are encouraged to be submitted electronically to <http://www.bogazicitipdergisi.com>. Along with revised form, the accepted paper in its final form also should be sent to www.bogazicitipdergisi.com.

BOSPHORUS MEDICAL JOURNAL

Boğaziçi Tıp Dergisi

SUBMISSION OF THE MANUSCRIPTS

Authors are required to submit the following:

- Copyright Transfer Form,
- Author Contributions Form.
- These forms are available for download at www.bogazicitipdergisi.com.

All correspondence will be sent to the first-named author unless otherwise specified. Corresponding author is expected to provide an e-mail address for all further correspondence. Papers should be accompanied by a cover letter indicating that the paper is intended for publication and specifying for which section of the Journal it is being submitted (i.e., original article, review article, or case report etc). Authors will be notified of the receipt of their paper and the number assigned to it. The number should be included in all further correspondence.

Authors should note, however, that manuscripts may be returned after initial review by the editorial office if the paper is deemed unlikely to be reviewed favorably by virtue of size restrictions and/or general interest for the readership. This rapid rejection process enables the author to promptly submit for publication elsewhere.

PREPARATION OF THE MANUSCRIPT

Maximum word length, number of authors, references, and table/figures should be based on the following table.

	Word	Reference	Table/Figure	Author
Research Article	6000	50	10	All
Review	6000	50	8	3
Case Report	2500	15	5	5
Editorial	2000	10	5	1
Letter to editor	500	5	2	2

General Format: All manuscripts should be typed using the standard A4-size format document with 2.5 cm-wide margins on all sides. The references should be numbered consecutively in the order of their first mention in the text. All text material, including references, footnotes, and table and figure legends, should be typed using double-spacing in an 11 point font with left alignment and without hyphenated line breaks. The fonts Times New Roman or Arial should be used in the text, for symbols, and all other special characters. Please use the editing features of your word processing program to type bold or italic letters, mathematical symbols, Greek letters, subscript and superscript characters. Please take care not to confuse the letters O and I with the numerals 0 and 1. To set a left indent for a paragraph, click the TAB button once. Only the International System of Units (SI) should be used for units of measurement. Please review the final version of the manuscript very carefully, especially for formatting and editing errors. Please note that American English spelling and terminology should be used in the manuscripts.

All pages of the manuscript should be consecutively numbered starting from the title page (page 1, title page; page 2, Turkish abstract (Turkish authors only); page 3, English abstract, etc.). Page numbers should be indicated on the upper right-hand corner of each page. The final electronic version of the manuscript should be in ".doc", ".docx" or ".rtf" format.

Title page: A separate title page should be submitted with all manuscripts and this page should include:

The full title of the manuscript as well as a short title (running head) of no more than 50 characters,

Name, affiliation(s), ORCID ID, and highest academic degree(s) of the author(s),
Details of any grant or other sources of support,

Name, address, telephone (including mobile phone number) and fax numbers, and email address of the corresponding author,

Acknowledgment of the individuals who contributed to the preparation of the manuscript but who do not fulfill authorship criteria.

Abstract: An abstract should be submitted with all manuscripts with the excep-

tion of Letters to the Editor. The abstract of an Original Article should be structured with subheadings (Objective, Methods, Results, and Conclusion). All acronyms and abbreviations used in the manuscript should be defined at first use, both in the abstract and in the main text. The abbreviation should be provided in parentheses following the definition.

Keywords: Each submission must be accompanied by a minimum of three to a maximum of six keywords at the end of the abstract to be used for subject indexing. The keywords should be listed in full without abbreviations. The keywords should be selected from the National Library of Medicine, Medical Subject Headings database (<https://www.nlm.nih.gov/mesh/MBrowser.html>).

Introduction: State the specific purpose and available data relevant to the study.

Methods: All methods used to select participants and conduct the study should be described in detail. Known methods should be cited. Novel or modified methods used should be described in detail. Doses, concentrations, routes, and duration of administration of drugs and chemical agents should be indicated. A concise report of all statistical methods used for summarizing available data and for testing the proposed hypothesis should be provided under a subtitle, including the p value criteria determined for statistically significant difference.

Statistical evaluation conducted should be explained in detail. Standard statistical methods should be used as much as possible. If rarely employed or novel statistical methods were used, then the relevant references should be cited.

When necessary, more detailed explanations about unusual, complex, or new statistical methods can be provided in separate files for readers as online supplementary data. When a trademarked drug, product, hardware, or software program is mentioned within the main text, product information, include the name of the product, the manufacturer of the product, and the city and the country of the company headquarters (including the state if in USA), should be provided in parentheses in the following format: "Discovery St PET/CT scanner (General Electric, Milwaukee, WI, USA). The recommendations in the statistics section of the "Uniform Requirements for Manuscripts Submitted to Biomedical Journals: Writing and Editing for Biomedical Publication" (<http://www.ICMJE.org>) should be taken into consideration.

Results: The study results should be presented in logical sequence and in detail. The findings should be supported by figures and tables. Information given in figures and tables should not be repeated in the text unless absolutely required.

Discussion: Data relevant to the study subject matter should be examined, evaluated, and substantiated with references from domestic and international sources. General information irrelevant or superfluous to the report should not be included.

Acknowledgement: The names of individuals who contributed to the study but who fail to meet the criteria of authorship should be mentioned in this section. The written consent of all individuals mentioned should be obtained.

The names of the authors and institutions should not be mentioned in the abstract, full text and table/figure legends. Author and institution information should only be stated on the title page. In addition, the names of the authors and institutions should not be included in the responses sent to the referees during the revision phase.

Manuscript Types

Boğaziçi Tıp Dergisi / Bosphorus Medical Journal publishes the types of articles briefly described below.

Research Articles: This is the most important type of article, since it provides new information based on original research. The main text of original articles should be structured with an Introduction, Methods, Results, Discussion, Conclusion, and References subheadings. Please see Table 1 for limitations for Research Articles.

Statistical analysis is usually necessary to support conclusions. Statistical analyses must be conducted in accordance with international statistical reporting standards (Altman DG, Gore SM, Gardner MJ, Pocock SJ. Statistical guidelines for contributors to medical journals. *Br Med J* 1983;7;1489-93). Information on statistical analyses should be provided with a separate subheading under the Materials and Methods section and the statistical software that was used during

BOSPHORUS MEDICAL JOURNAL

Boğaziçi Tıp Dergisi

the process must be specified.

Units should be prepared in accordance with the International System of Units (SI).

Limitations, drawbacks, and the shortcomings of original articles should be mentioned in the Discussion section before the conclusion paragraph.

Review Articles: Reviews prepared by authors who have extensive knowledge of a particular field and whose scientific background has been translated into a large volume of publications with a high citation potential are welcomed. Submissions from such authors may also be invited by the journal. Reviews should describe, discuss, and evaluate the current level of knowledge of a topic in clinical practice and should guide future studies. The main text should include an Introduction, Clinical and Research Consequences, and Conclusion sections. Please refer to Table 1 for the limitations for Review Articles.

Case Reports: There is limited space for case reports in the journal and reports on rare cases or conditions that constitute challenges in diagnosis and treatment, those offering new therapies or revealing knowledge not included in the literature, and interesting and educative case reports are accepted for publication. The text should include an Introduction, Case Presentation, and Discussion subheadings. Interesting and unusual images are an advantage in the evaluation process. Please see Table 1 for the limitations for Case Reports.

Letters to the Editor: This type of manuscript discusses important parts, overlooked aspects, or lacking parts of a previously published article. Articles on subjects within the scope of the journal that might attract the readers' attention, particularly educative cases, may also be submitted in the form of a Letter to the Editor. Readers can also present their comments on published manuscripts in the form of a Letter to the Editor. An abstract, Keywords, and Tables, Figures, Images, and other media should not be included. The text should not include subheadings. The manuscript that is being commented on must be properly cited within this manuscript.

References

References are numbered and listed by their order of appearance in text; the text citation is followed by the appropriate reference number in parentheses. References should be restricted to closely pertinent material. Accuracy of citation is the author's responsibility. References should conform exactly to the original spelling, accents, punctuation, etc. All references should be cited inside the text. References must be listed in the Vancouver Style only. All references should be numbered sequentially as superscripted in the text and listed in the same numerical order in the reference section. The reference styles for different types of publications are presented in the following examples.

JOURNAL REFERENCE

Uruc F, Yildirim C, Aras B, Kivrak M, Sahin A, Akan S, et al. Infectious Complications of Transrectal Ultrasound-guided Prostate Needle Biopsy: A Retrospective Case-Control Study. *Bosphorus Med J* 2016; 3:5-8.

Journal Article:

Matsukawa Y, Kato K, Hatta Y, Iwamoto M, Mizuno S, Kurihara R, et al. Helicobacter pylori eradication reduces platelet count in patients without idiopathic thrombocytopenic purpura. *Platelets* 2007;18:52-5.

Book Chapter Reference

Author. Title. In: Editor, 'Ed.'.^'Eds.'. Book Title. ed. Edition, Place Published: Publisher; Year: p. Pages. Lal G, Clark OH. Thyroid, parathyroid, and adrenal. In: Brunicaudi FC, Andersen DK, Billiar TR, Dunn DL, Hunter JG, Matthews JB, Eds. Schwartz's Principles of Surgery. ed.10th, New York; Mc-Graw Hill; 2015: p.1521-96.

Book Reference

Author. Title. ed. Edition, Place Published: Publisher; Year. Walsh P. Physiology and Pharmacology of the Bladder and Urethra. *Campbell's Urology*. ed. 10th, Philadelphia: Saunders; 2014.

Conference Proceedings:

Author. Title. In: Editor, 'editor'.^'editors'. Conference Name; Year of Conference Date; Conference Location: Publisher; Year of Conference]. p. Pages.

Bengissson S. Sothemin BG. Enforcement of data protection, privacy and security in medical informatics. In: Lun KC, Degoulet P, Piemme TE, Rienhoff O, editors. *MED-INFO 92. Proceedings of the 7th World Congress on Medical Informatics*; 1992 Sept 6-10; Geneva, Switzerland. Amsterdam: North-Holland; 1992. pp.1561-5.

Scientific or Technical Report:

Author. Title. Type. Place Published: Institution; Year Date. Report No.: Report Number.

Cusick M, Chew EY, Hoogwerf B, Agrón E, Wu L, Lindley A, et al. Early Treatment Diabetic Retinopathy Study Research Group. Risk factors for renal replacement therapy in the Early Treatment Diabetic Retinopathy Study (ETDRS), Early Treatment Diabetic Retinopathy Study Kidney Int: 2004. Report No: 26.

Thesis:

Author. Title. Type. Place Published: Institution; Year Date. Report No.: Report Number.

Kaplan SI. Post-hospital home health care: elderly access and utilization (dissertation). St Louis (MO): Washington Univ; 1995.

Epub Ahead of Print Articles:

Author. Title. Alternate Title Year Date Accessed. doi: DOI. [Epub ahead of print].

Cai L, Yeh BM, Westphalen AC, Roberts JP, Wang ZJ. Adult living donor liver imaging. *Diagn Interv Radiol*. 2016 Feb 24;doi: 10.5152/dir.2016.15323. [Epub ahead of print].

URL (WebPage)

Author. Title. Available from: URL. Accessed Access Date, Access Year. Fox S. Pew Research Center. The social life of health information. 2014. Available from: <http://www.pewresearch.com>.

PUBLISHING FEE

There is no fee for article submission, article processing or publication. Manuscripts can be reached from the web page of journal without any fees. No additional fee is required from the authors for accepted manuscripts.

OPEN ACCESS AND COMMONS USER LICENSES

Open Access

Boğaziçi Tıp Dergisi / Bosphorus Medical is an open access journal which means that all content is freely available without charge to the user or his/her institution. Users are allowed to read, download, copy, distribute, print, search, or link to the full texts of the articles, or use them for any other lawful purpose, without asking prior permission from the publisher or the author. This is in accordance with the BOAI definition of open access.

Commons User Licenses

This work is licensed under a Creative Commons Attribution-Non Commercial 4.0 International License (CC BY-NC 4.0) .

Plagiarism Detection:

Journal editors and readers have a right to expect that a submitted work is the author's own contribution that has not been plagiarized (i.e. taken from other authors without permission, if permission is required) and that copyright has not been breached (for example, if figures or tables are reproduced). The editorial office runs a plagiarism check on every submission using iThenticate. By submitting your manuscript to this journal you accept that your manuscript may be screened for plagiarism against previously published works. Individual authors and researchers can now check their work for plagiarism before submission.

Identification of and dealing with allegations of research misconduct

The publisher and the editor of the journal take reasonable steps to identify and prevent the publication of papers where research misconduct has occurred, including plagiarism, under/over citation, and data falsification/fabrication, among others. In no case shall the journal or its editor encourage such misconduct, or knowingly allow such misconduct to take place. In the event that the journal's publisher or editor is made aware of any allegation of research misconduct relating to a published article in the journal, the publisher or editor shall follow COPE's guidelines (or equivalent) in dealing with allegations.

OPEN ACCESS POLICY

Bogazici Tıp Dergisi (Bosphorus Medical Journal) supports the Budapest Open Access Initiative statement of principles that promotes free access to research literature. The declaration defines open access to academic literature as free availability on the internet, permitting users to read, record, copy, print, search, or link to the full text, examine them for indexing, use them as data for software or other lawful purposes without financial, legal, or technical barriers. Information sharing represents a public good, and is essential to the advancement of science. Therefore, articles published in this journal are available for use by researchers and other readers without permission from the author or the publisher provided that the author and the original source are cited. The articles in **Bogazici Tıp Dergisi (Bosphorus Medical Journal)** are accessible through search engines, websites, blogs, and other digital platforms.

Additional details on the Budapest Open Access Initiative and their guidelines are available at <https://www.budapestopenaccessinitiative.org/>, including a Turkish translation of the recommendations at <http://www.budapestopenaccessinitiative.org/boai-10-translations/turkish-translation>.

FEE - CHARGES

This journal assesses no submission fees, publication fees (article processing charges), or page charges.

CREATIVE COMMONS LICENSE

A Creative Commons license is a public copyright license that enables the free distribution of copyrighted work. **Bogazici Tıp Dergisi (Bosphorus Medical Journal)** articles are licensed under the Attribution-NonCommercial-ShareAlike 4.0 International (CC BY-NC 4.0) version. The author grants the right to share and use original work with the condition that it be appropriately credited, it may not be used for commercial purposes, and secondary products must also be made available under the same terms of use. Specific details can be found at <https://creativecommons.org/licenses/by-nc/4.0/>.

Bogazici Tıp Dergisi (Bosphorus Medical Journal) is committed to encouraging dissemination of academic work and interdisciplinary cooperation.

ETHICS POLICY

The observance of ethical principles throughout the research and publication process is fundamental to ensuring the integrity of the work and furthering the goal of contributing to and sharing high-quality, objective, reliable, and useful information.

Bogazici Tıp Dergisi (Bosphorus Medical Journal) has adopted ethical principles based on the guidelines prepared by the Committee on Publication Ethics (COPE) (<https://publicationethics.org/>). We implement these processes to ensure appropriate support for our authors and their institutions, as well as our readers. It is crucial that all of the stakeholders in the process (authors, readers and researchers, publishers, reviewers, and editors) comply with ethical principles.

ARTIFICIAL INTELLIGENCE (AI)-ASSISTED TECHNOLOGY

At submission, the journal should require authors to disclose whether they used artificial intelligence (AI)-assisted technologies (such as Large Language Models [LLMs], chatbots, or image creators) in the production of submitted work. Authors who use such technology should describe, in both the cover letter and the submitted work, how they used it. Chatbots (such as ChatGPT) should not be listed as authors because they cannot be responsible for the accuracy, integrity, and originality of the work, and these responsibilities are required for authorship. Therefore, humans are responsible for any submitted material that included the use of AI-assisted technologies. Authors should carefully review and edit the result because AI can generate authoritative-sounding output that can be incorrect, incomplete, or biased. Authors should not list AI and AI-assisted technologies as an author or co-author, nor cite AI as an author. Authors should be able to assert that there is no plagiarism in their paper, including in text and images produced by the AI. Humans must ensure there is appropriate attribution of all quoted material, including full citations.

ETHICAL RESPONSIBILITIES OF THE AUTHORS

- Studies submitted for publication must be original work of the author. References to other studies must be cited and/or quoted completely and accurately;
- Only those who provide a substantial intellectual contribution to the content of the work may be cited as an author. Other contributors may be recognized with acknowledgements at the conclusion of the article;
- All competing interests or relationships that may be perceived to constitute a conflict of interest must be declared and explained;
- All studies involving human or animal subjects must comply with national and international laws and guidelines regarding privacy and ethical conduct (e.g., World Medical Association Declaration of Helsinki, US National Institutes of Health Policy on the Use of Laboratory Animals, EU Directive on the Use of Animals) and the details of approval and observance should be indicated in the Materials and Methods section of the manuscript;
- Authors must be able to provide documentation showing that they have the right to use the data analyzed, all necessary permission related to the research, and appropriate consent;
- Raw data and other material used in the article must be available and may be requested from the author(s) in order to verify the validity of the reporting;
- In the event the author(s) notice an error at any point in the publication process or after publication, they have the obligation to inform the journal editor or publisher and cooperate in appropriate corrective action;
- Authors may not submit their article for publication to more than one journal simultaneously. Each application must be initiated following the completion of any previous effort. Previously published articles, will not be accepted, including translations, without the proper acknowledgement of the original author;
- Changes in authorship designation (such as adding authors, changing the printed order of the authors, removing an author) once the evaluation process has begun will not be accepted in order to protect all parties involved.

ETHICAL DUTIES AND RESPONSIBILITIES OF THE EDITORS

The editor is responsible for everything published in the journal. In the context of this responsibility, editors have the following duties and obligations:

- Endeavor to meet the needs of readers and authors;
- Maintain continuous development to improve the journal;
- Consistently work to ensure quality and academic integrity. The editor is responsible for confirming that the publishing policies and standards are upheld;
- Support freedom of thought;
- Prevent business needs or other considerations from compromising intellectual and ethical standards, including acting in a balanced, objective, and fair manner in the course of their duties without any discrimination based on gender, religious or political beliefs, ethnic or geographical origin, sponsorship, renown, or other influence;
- Apply the publicly defined publication policies created and enforced to ensure a timely and impartial evaluation process for all submissions;
- Protect intellectual property and to defend the rights of the journal and author(s);
- Demonstrate clarity and transparency. The editor is expected to ensure that any errors, inconsistencies, or misleading statements are corrected quickly and appropriately acknowledged;
- Perform a thorough, timely, and objective investigation of any complaint or allegation of misconduct, including providing the opportunity for the author to present information refuting accusations, and to share the findings and conclusions and implement appropriate action, which may include, but is not limited to rejection of an article.

BOSPHORUS MEDICAL JOURNAL

Boğaziçi Tıp Dergisi

READER RELATIONSHIP

The editor is to make publication decisions based on expectations of suitable and desirable material. Studies accepted for publication must be original contributions that benefit the reader, researcher, practitioner, and the literature. In addition, editors are obliged to take into account feedback from readers, researchers, and practitioners, and to provide an informative response. Readers will also be informed of any funding provided to support published research.

AUTHOR RELATIONSHIP

- The decision to accept an article is to be based on the importance, original value, validity, and clarity of expression of the work, and the goals and objectives of the journal;
- Studies accepted for evaluation and publication will not be withdrawn unless serious problems are identified;
- The editor will not disregard positive reviewer comments unless there is a serious problem with the study;
- New editors will not change publishing decisions made by previous editor(s) unless there is a serious problem;
- A description of the submission and evaluation process is publicly available;
- Authors are provided with descriptive and informative feedback.

REVIEWER RELATIONSHIP

- Reviewers are to be selected according to the subject of the study;
- Information and guidance for the evaluation phase is provided;
- Any conflicts of interest between authors and reviewers will be disclosed and managed appropriately;
- Reviewer identity is to be kept confidential to preserve a blind review process;
- Reviewers are to evaluate the study using unbiased, scientific, and constructive comments. Unkind or unscientific commentary will not be permitted;
- Reviewers will be evaluated using criteria such as timely response and quality of observations;
- The pool of reviewers is to be assessed and supplemented regularly to ensure a broad scope of expertise.

EDITORIAL BOARD RELATIONSHIP

The editor works with the members of the editorial board to ensure that they are familiar with journal policies and developments in regular meetings and announcements, and will provide training for new members and assistance to board members during their tenure in their role as a supporter of the journal.

- Editorial board members must be qualified and able to contribute to the journal;
- Members of the editorial board must evaluate studies impartially and independently;
- Editorial board members with the appropriate expertise will be given the opportunity to evaluate suitable articles;
- The editor will maintain regular contact with the editorial board and hold regular meetings regarding the development of editorial policies and other aspects of journal management.

Creativity and Openness

- Constructive criticism is to be encouraged;
- Authors will be given the opportunity to reply to criticism or lodge complaints;
- Negative results will not be a reason for submission denial.

ETHICAL RESPONSIBILITIES OF THE REVIEWERS

Peer review of research embodies the scientific method, subjecting the work to the exacting scrutiny of knowledgeable colleagues. The rigor of the review process directly affects the quality of the literature; it provides confidence in an objective and independent evaluation of the published work. **Bogazici Tıp Dergisi**

(**Bosphorus Medical Journal**) uses a double-blind review process. All comments and the evaluation are transmitted through the journal management system. Reviewers should:

- Only agree to evaluate studies related to their specialty;
- Return reviews promptly and within the designated timeframe;
- Evaluate with impartiality. Nationality, gender, religious beliefs, political beliefs, commercial concerns, or other considerations must not influence the evaluation;
- Refuse to review any work with a potential conflict of interest and inform the journal editor;
- Maintain confidentiality of all information. Destroy manuscripts and related material following the review. Only the final published version may be used for any purpose;
- Report any suspicion of misconduct to the editor;
- Use thoughtful and constructive language intended to improve the quality of the article. Hostile or derogatory comments are not acceptable.

ETHICAL RESPONSIBILITIES OF THE PUBLISHER

The role of the publisher includes stewardship of the scholarly record. As such, the publisher should:

- Abide by ethical principles related to research integrity; the process of application, review, and selection; and publication;
- Collaborate with the editor and the editorial board to maintain and develop the journal in a relationship that recognizes editorial independence and is defined by written agreement;
- Publish content in a timely manner, including corrections, clarifications, and retractions;
- Preserve published work.

The publication processes of the journal are conducted in accordance with the guidelines of International Committee of Medical Journal Editors (ICMJE), the World Association of Medical Editors (WAME), the Council of Science Editors (CSE), the Committee of Publication Ethics (COPE), the European Association of Science Editors (EASE) and National Information Standards Organization (NISO).

Examples of some activities considered to be contrary to scientific research and publication ethics:

- **Plagiarism:** The representation of the ideas, methods, data, or other work of another, in whole or in part, as one's own. The original source must be appropriately acknowledged. Authors are encouraged to offer unique work that does not rely on substantial use of other sources, regardless of citation.
- **Fraud:** The use of fabricated or falsified data or other deceptive misrepresentation of fact.
- **Distortion:** Manipulation of the research records, data, images, or results, or presenting unused devices or materials as if they were used in the research, particularly in the interests of study sponsors.
- **Republication:** Duplicate submissions presented as unique publications.
- **Slicing:** The use of a portion of data or findings derived from a single research idea in multiple smaller units as separate publications.
- **Inaccurate authorship:** The inclusion of individuals as named authors who were not active contributors, the failure to include contributors, or the inappropriate ranking of authors.
- **Lack of acknowledgement** of individuals, institutions, or organizations that provided financial or other substantial support to the work.
- **Use of a thesis or unpublished study** without the permission of the owner.
- **Failure to comply with ethical rules** for human and animal research, including respect for patient rights and animal welfare, or obtaining the required approval.
- **The misuse of resources, facilities, or devices** provided for scientific research.

- The use of false or misleading statements.

On rare occasions it may be necessary for a journal to impose sanctions on researchers who have engaged in questionable research practices or publishing ethics malpractice: for example, a ban against publishing any further articles in the journal when doing so puts the journal's reputation demonstrably at risk, or not permitting a researcher to serve as a reviewer or editor. Bans of this nature may be implemented for a period of time and revoked or extended if necessary, upon review at the conclusion of the allotted time period. Sanctions may be appealed by writing to the journal editor.

PLAGIARISM POLICY

Plagiarism is the theft of another's work and a violation of ethics, regardless of whether it is intentional or not. It is unacceptable conduct to submit or publish manuscripts using other sources without appropriately citing the reference. It is the policy of the **Bogazici Tıp Dergisi (Bosphorus Medical Journal)**, to use plagiarism detection software for all submissions and to perform an editorial review when necessary. The editor or the editorial board may request revision or reject a manuscript that does not meet publication standards, including plagiarism, citation or other manipulation, or any fraudulent misrepresentation.

COPYRIGHT TRANSFER

Manuscripts submitted for publication in the **Bogazici Tıp Dergisi (Bosphorus Medical Journal)** should be original, unpublished work. Upon submission, the authors are obliged to declare that the study, in whole or in part, has not been previously published or evaluated for publication on any other platform. Sanctions may be applied for failure to observe this policy.

Authors agree to transfer copyright privileges upon submission to the **Bogazici Tıp Dergisi (Bosphorus Medical Journal)**. This transfer takes effect upon acceptance for publication. No part of published material may be used for any other purpose without the written permission of the publisher.

Authors should obtain any necessary permission from the copyright holder

when using content previously published in printed or electronic format, including pictures, tables, or other elements. The legal, financial, and criminal responsibility resides with the author.

Authors must return a completed copyright transfer form upon submission.

CONFLICT OF INTEREST

The editor is required to ensure that any conflicts of interest between authors, reviewers, or other parties are disclosed and managed appropriately to provide an independent and impartial process.

Any potential perception of a financial or personal interest that may affect decision-making creates a conflict of interest. The presence of a conflict of interest is independent of the occurrence of inappropriateness. The reliability of the scientific process and published articles is directly related to the objective consideration of conflicts of interest during the planning, implementation, writing, evaluation, editing, and publication of scientific studies.

Financial relationships are the most easily identified conflicts of interest, and if undisclosed, they undermine the credibility of the journal, the authors, and the science. However, conflicts can also occur through individual relationships, academic competition, intellectual approach, and more. Authors should refrain as much as possible from any relationship that could restrict their ability to objectively access data or analyze, interpret, prepare, and publish their article. Authors must disclose any relationships related to study submissions.

Editors and peer reviewers should also be aware of potential conflicts of interest and refrain from engaging in any activity that could be questionable and report associations that could be perceived as presenting a conflict.

The publication team works diligently to ensure that the evaluation process is conducted in an impartial manner in order to protect the interests of all parties.

The conflict of interest form and more detailed information are available at: <http://www.icmje.org/disclosure-of-interest/>

ORIGINAL ARTICLES

- Comparison of Functional and Pain Scores in The Early Period in Patients Undergoing Arthroscopic Rotator Cuff Repair Under General Anesthesia or Interscalene Block..... 27**
Genel Anestezi veya İnterskalen Blok Altında Artroskopik Rotator Manşet Onarımı Yapılan Hastalarda Erken Dönemde Fonksiyonel ve Ağrı Skorlarının Karşılaştırılması
Çakar A, Karakuş Ö, Yılmaz B
- Assessment of Determinants Influencing Fall Risk Among Geriatric Patients 38**
Geriatrik Hastalarda Düşme Riskini Etkileyen Faktörlerin Değerlendirilmesi
Yorulmaz E, Günay B, Mesci N, Üstün B, Geler Külcü D
- Performance of the ‘Turkish Triage System’ During the Covid-19 Pandemic 48**
COVID-19 Pandemisinde ‘Türk Triage Sistemi’nin Performansı
Akyıldız SF, Sarı Doğan F, Cimilli Öztürk T, Ünal Akoğlu E, Bostancı Y
- Comparison of the Efficacy of Low-Level Laser Therapy and Ultrasound Treatment in Patients with Shoulder Pain 56**
Omuz Ağrılı Hastalarda Düşük Yoğunluklu Lazer ve Ultrason Tedavisinin Etkinliğinin Karşılaştırılması
Aydeniz B, Uslu T
- Effects of Greater Occipital Nerve Block on Headache Frequency and Sleep Disturbance in Medication Overuse Headache .. 63**
Büyük Oksipital Sinir Bloğunun İlaç Aşırı Kullanım Baş Ağrısında Baş Ağrısı Sıklığı ve Uyku Bozukluğu Üzerine Etkileri
Akpınar S.

CASE REPORT

- An Early-Stage Neuroborreliosis Case with Atypical Findings 69**
Atipik Bulgularıyla Erken Dönem Nöroborreliyoiz Olgusu
Yapıcı S, Söylemez E, Özdağ MF, Karşıdağ S.



Comparison of Functional and Pain Scores in The Early Period in Patients Undergoing Arthroscopic Rotator Cuff Repair Under General Anesthesia or Interscalene Block

Genel Anestezi veya İnterskalen Blok Altında Artroskopik Rotator Manşet Onarımı Yapılan Hastalarda Erken Dönemde Fonksiyonel ve Ağrı Skorlarının Karşılaştırılması

¹Department of Orthopedics and Traumatology, Balıkesir Atatürk City Hospital, Balıkesir, Türkiye

²Department of Orthopedics and Traumatology, University of Health Sciences, Balıkesir Atatürk City Hospital, Balıkesir, Türkiye

³Department of Orthopedics and Traumatology, University of Health Sciences, Fatih Sultan Mehmet Training and Research Hospital, Istanbul, Türkiye

Arif Çakar,¹ Özgün Karakuş,² Barış Yılmaz³

ABSTRACT

Objectives: This study aimed to compare the effects of general anesthesia (GA) and interscalene brachial plexus block (ISB) on clinical outcomes following arthroscopic single-row or double-row repair of crescentic rotator cuff tears.

Methods: This retrospective cohort study included 80 patients (46 women, 34 men; mean age, 66.7 years) who underwent arthroscopic repair for crescent-shaped full-thickness rotator cuff tears. Patients were divided into GA (n=40) and ISB (n=40) groups. Tears were classified according to the Cofield classification. Functional outcomes were assessed using Constant-Murley, ASES, and UCLA scores, along with joint range of motion (JROM) and visual analog scale (VAS) pain scores. Evaluations were performed at 6 and 18 months postoperatively.

Results: At 6 months, Constant-Murley, ASES, and UCLA scores were significantly higher in the ISB group than in the GA group. No significant differences were observed between the groups at 18 months. Both groups showed significant improvement over time. JROM and VAS scores were similar between the groups. Demographic variables and repair type had no significant effect on outcomes.

Conclusion: ISB was associated with improved short-term functional outcomes following arthroscopic rotator cuff repair. Long-term clinical results were comparable to those achieved with general anesthesia.

Keywords: ASES; Constant-Murley score; Crescent-shaped rotator cuff tear; General anesthesia; Interscalene block; UCLA score.

ÖZET

Amaç: Bu çalışma, genel anestezi (GA) ve interskalen brakial pleksus bloğu (ISB) uygulamalarının, artroskopik tek sıra veya çift sıra hilal şeklindeki rotator manşet yırtıklarının onarımından sonraki klinik sonuçlar üzerindeki etkilerini karşılaştırmayı amaçlamıştır.

Yöntem: Bu retrospektif kohort çalışması, hilal şeklinde tam kat rotator manşet yırtıkları nedeniyle artroskopik onarım uygulanan 80 hastayı (46 kadın, 34 erkek; ortalama yaş 66,7) içermektedir. Hastalar GA (n=40) ve ISB (n=40) gruplarına ayrılmıştır. Yırtıklar Cofield sınıflandırmasına göre sınıflandırılmıştır. Fonksiyonel sonuçlar, Constant-Murley, ASES ve UCLA skorları ile eklem hareket açıklığı (JROM) ve görsel analog skala (VAS) ağrı skorları kullanılarak değerlendirilmiştir. Değerlendirmeler ameliyat sonrası 6. ve 18. aylarda yapılmıştır.

Cite this article as:

Çakar A, Karakuş Ö, Yılmaz B. Comparison of Functional and Pain Scores in The Early Period in Patients Undergoing Arthroscopic Rotator Cuff Repair Under General Anesthesia or Interscalene Block Bosphorus Med J 2026;13(2):27-37.

Received: 09.10.2025

Revision: 23.12.2025

Accepted: 09.09.2026

Correspondence:

Dr. Arif Çakar.
Department of Orthopedics and Traumatology, Balıkesir Atatürk City Hospital, Balıkesir, Türkiye

Phone:

+90 536 879 78 18

e-mail:

arifcakar94@gmail.com

OPEN ACCESS



This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License.

Bulgular: Altıncı ayda Constant-Murley, ASES ve UCLA skorları, GA grubuna kıyasla ISB grubunda anlamlı olarak daha yüksekti. On sekizinci ayda gruplar arasında anlamlı bir fark gözlenmedi. Her iki grup da zaman içinde anlamlı bir iyileşme gösterdi. JROM ve VAS skorları gruplar arasında benzerdi. Demografik değişkenler ve onarım tipi, sonuçlar üzerinde anlamlı bir etkiye sahip değildi.

Sonuç: ISB, artroskopik rotator manşet onarımı sonrası kısa dönem fonksiyonel sonuçlarda iyileşme ile ilişkiliydi. Uzun dönem klinik sonuçlar, genel anestezi ile elde edilen sonuçlarla karşılaştırılabilir.

Anahtar sözcükler: ASES; Constant Murley Skoru; Genel Anestezi; Interskalen blok; Kresent tip rotator cuff yırtığı; UCLA skoru.

Conservative treatment is the first choice for rotator cuff injuries, especially in cases of partial tears and in patients who are not suitable for surgery. The aim is to control pain, improve function, and prevent progression of the tear. Response to treatment usually occurs within 6–12 weeks.^[1-3] Surgical treatment involves open, mini-open, and arthroscopic repair techniques. In full-thickness tears, the tendon is completely detached from the humeral footprint, and repair is performed using one of these three methods.^[4-6]

Arthroscopic rotator cuff repair is frequently preferred today because of both technical advances and its less invasive nature. High success rates have been reported, especially for small and medium-sized tears. The advantages of this method include minimal soft tissue damage, rapid functional recovery, and less postoperative pain.^[6]

General anesthesia is the standard method in shoulder surgery because it ensures reliable airway control; however, it is more prone to hemodynamic fluctuations such as hypotension and bradycardia. While central blocks require caution in patients using anticoagulants, peripheral nerve blocks offer safe alternatives. The interscalene brachial plexus block (ISB) is the preferred regional anesthesia method in shoulder surgery because it provides effective motor and sensory blockade. However, ISB administration requires an experienced anesthesiologist and appropriate patient preparation.^[7,8]

The aim of our study was to evaluate the use of general anesthesia and interscalene block anesthesia methods for the arthroscopic repair of crescentic (crescent-shaped) rotator cuff tears in patients undergoing single- or double-row repair and, additionally, to evaluate functional shoulder scores, shoulder joint range of motion, and pain scales according to tear size, repair method, affected shoulder, and sociodemographic characteristics (age, sex, and body mass index [BMI]).

There is insufficient information in the literature regarding functional differences between ISB and GA techniques in

patients undergoing rotator cuff repair in the short and medium term. In our study, we expect better results in terms of functional and pain scales with the ISB technique in the short term, while we anticipate equivalent results in the medium term.

Methods

Ethical approval was obtained from the Scientific Research Ethics Committee of Balıkesir Atatürk City Hospital on 19.06.2025, with decision no. 2025/06/60. Informed consent was waived for this study, which was conducted in accordance with the Declaration of Helsinki.

This retrospective cohort study included a total of 80 patients who underwent single- or double-row rotator cuff repair using arthroscopic techniques. All cases were evaluated with preoperative magnetic resonance imaging (MRI) and plain radiographs; only patients with a crescentic full-thickness tear were included in the study. The morphology and size of the tears were defined intraoperatively by the surgeon and classified according to the Cofield classification as small (<1 cm), medium (1–3 cm), large (3–5 cm), and massive (>5 cm), based on the surgical reports. All surgical reports were retrospectively reviewed by two orthopedic surgeons.

Patients were divided into two groups with equal numbers (n=40).

Group A included patients operated on under general anesthesia, with 6 patients having small tears, 26 patients having medium tears, 7 patients having large tears, and 1 patient having a massive tear. Group B consisted of patients who underwent anesthesia with an interscalene brachial plexus block (ISB), including 6 patients with small tears, 26 patients with medium tears, 7 patients with large tears, and 1 patient with a massive tear.

In both groups, a single-row repair technique was applied to 20 patients and a double-row repair technique to 20 patients. Acromial spurs identified during arthroscopy were minimally shaved using a burr.

Patients with the following characteristics were excluded from the study: shoulder instability, degenerative arthritis of the glenohumeral joint, acromioclavicular joint pathology, tubercle fracture, previous shoulder surgery, revision surgery due to re-rupture, massive contracted and immobile tears, cases requiring subscapularis repair, and cases with Goutallier stage IV fatty degeneration.

Patients were followed for an average of 24.2 months (range, 18–32 months). During follow-up, all patients were re-evaluated at the 6- and 18-month follow-up visits, including assessments of functional status (ROM and functional shoulder scores) and pain (VAS score), along with any additional medical conditions and body mass index. Since the shortest follow-up period was 18 months, the 18-month follow-up was used for the final evaluation of all patients in the study.

The functional assessment scales applied included the Constant-Murley Score (CMS), American Shoulder and Elbow Surgeons (ASES) score, University of California at Los Angeles (UCLA) score, and range of motion (ROM) measurements. ROM values were measured using a goniometer in the directions of flexion, abduction, internal rotation, and external rotation. All measurements were performed in a standardized manner by two orthopedic surgeons during the patients' 6- and 18-month follow-up visits.

Surgical Technique

All procedures were performed with the patients in the supine position. First, arthroscopic examination of the glenohumeral joint was performed via the posterior portal. Subsequently, the subacromial space was accessed to visualize the rotator cuff tear. Next, the footprint was identified, and using the lateral portal, a sliding knot was tied with a 4.5-mm titanium anchor (Artrrotek®) from SMC (Samsung Medical Center), compressing the rotator cuff and completing the single-row repair. Two medial anchors were used for tears that could not be completely closed. In addition, some cases were repaired in two rows using a 4.5-mm PushLock anchor with the tension-band method. Subacromial decompression was performed in all cases.

For the ultrasound-guided interscalene block, the linear probe was placed transversely in the supraclavicular fossa and advanced cranially to visualize the short axis of the brachial plexus, identifying the C5–C6 roots between the scalene muscles. The dorsal scapular and long thoracic nerves, which run within the middle scalene muscle, were

identified and preserved. Using an in-plane technique, an 18-G Tuohy needle was directed laterally toward the plexus sheath with the aid of hydrodissection. After ruling out intravascular injection by negative aspiration, 20 mL of 0.25% bupivacaine was administered in fractional 5-mL injections. The procedure was completed with a sterile dressing containing chlorhexidine gluconate.

The postoperative follow-up protocol was the same for all groups. Passive shoulder movement was initiated 1 day after surgery in all patients. A shoulder-arm sling with abduction support was used for 6 weeks, and active shoulder movement was restricted during this period. After 6 weeks, the shoulder-arm sling was removed, and all patients underwent a 2-month physical therapy program for active shoulder movements. No complications were observed in any patient during postoperative follow-up.

Statistical Analyses

Data analysis was performed using SPSS version 27.0 (IBM Corp., Armonk, NY, USA). Descriptive statistical methods (mean, standard deviation, minimum, and maximum) were used to evaluate the study data. The normality of quantitative data was tested using the Shapiro-Wilk test and graphical evaluations. For parametric data, the independent-samples t-test and paired t-test were used for intergroup comparisons, the mixed ANOVA test was used for repeated measures, and the chi-square test was used to compare categorical variables. The significance level was set at $p < 0.05$.

Results

A retrospective cohort study was conducted, and 80 patients with complete crescentic rotator cuff tears who underwent single-row or double-row arthroscopic repair between January 2018 and January 2025 were randomly selected for inclusion according to the inclusion criteria. Sixty-one point two five percent ($n=49$) of the patients were female, and thirty-eight point seven five percent ($n=31$) were male, with a mean age of 64.29 years (range, 50–81 years) (Table 1).

Eighty patients were randomly assigned to two groups for evaluation. Group A received general anesthesia, while Group B received an interscalene brachial plexus block (ISB).

In Group A, there were 40 patients in total, 16 of whom were male and 24 were female. The youngest patient was 50 years old and the oldest was 81 years old, with a mean age of 64.29

Table 1. Demographic data of patients

	Group A (General Anesthesia)	Group B (Interscalenous Brachial Plexus Block)	Total
Number of patients	40	40	80
Gender (Male-Female)	16-24	15-25	80
Age	Mean: 64.29 (Range: 50–81 years)	Mean: 64.28 (Range: 50–81 years)	Mean: 64.29 (Range: 50–81 years)
Side			
Right	25	24	49
Left	15	16	31
Torn Type			
Small	6	6	12
Medium	26	26	52
Large	7	7	14
Massive	1	1	2
Follow-up period	Mean: 24.2 months (Range: 18–32 months)	Mean: 24.2 months (Range: 18–32 months)	Mean: 24.2 months (Range: 18–32 months)
Single row repair	20	20	40
Double row repair	20	20	40

years. Twenty-five patients had a crescent-type rotator cuff tear in the right shoulder, and 15 patients had one in the left shoulder.

At the 6-month follow-up, shoulder flexion was 150–162 degrees (mean, 157.3), abduction was 152–160 degrees (mean, 156.8), internal rotation was 52–59 degrees (mean, 55.6), and external rotation was 57–63 degrees (mean, 59.7). At the 18-month follow-up, shoulder flexion was 150–162 degrees (mean, 157.6), abduction was 152–159 degrees (mean, 156.4), internal rotation was 50–58 degrees (mean, 55.5), and external rotation was 57–63 degrees (mean, 59.6), as measured by goniometry (Table 2).

Repair was performed using a single-row technique in 20 patients and a double-row technique in 20 patients. Tear size was reported as small in 6 patients, medium in 26 patients, large in 7 patients, and massive in 1 patient.

Patients' functional scores at 6 months postoperatively showed a Constant-Murley score of 76–85 (mean, 80.23), an ASES score of 74–88 (mean, 80.45), a UCLA score of 24–32 (mean, 27.50), and a VAS score of 0–3 (mean, 1.14). At 18 months postoperatively, the Constant-Murley score was 77–86 (mean, 80.95), the ASES score was 76–89 (mean, 81.88), the UCLA score was 24–32 (mean, 27.67), and the VAS score was 0–3 (mean, 1.12) (Table 3).

Table 2. Joint range of motion values for patients in groups at 6 and 18 months

	Group A (General Anesthesia)		Group B (Interscalene Brachial Plexus Block)	
	6 th month Min-max (Mean) (SD)	18 th month Min-max (Mean) (SD)	6 th month Min-max (Mean) (SD)	18 th month Min-max (Mean) (SD)
Shoulder flexion degree	150-162 ° (Mean:157.3°) (3.5)	150-162 ° (Mean:157.6°) (3.3)	153-165 ° (Mean:158.1°) (3.3)	153-164 ° (Mean:157.9°) (3.3)
Shoulder abduction degree	152-160 ° (Mean:156.8°) (2.2)	152-159 ° (Mean:156.4°) (2.2)	152-160 ° (Mean:157.2°) (2.2)	151-158 ° (Mean:156.5°) (2.2)
Degree of shoulder internal rotation	52-59 ° (Mean:55.6°) (2.1)	50-58 ° (Mean:55.5°) (2.3)	52-60° (Mean:55.8°) (2.0)	51-59° (Mean:55.7°) (2.0)
Shoulder external rotation degree	57-63 ° (Mean:59.7°) (2.1)	57-63 ° (Mean:59.6°) (2.1)	56-63° (Mean:59.8°) (2.0)	55-63° (Mean:59.7°) (2.1)

SD: Standard deviation.

Table 3. Postoperative shoulder function scores in patients undergoing surgery under general anesthesia (GA)

Functional scores and pain scales	6 th month Min-Max (Mean)	SD (6 th month)	18 th month Min-Max (Mean)	SD (18 th month)
Constant Murley Score	76–85 (80.22)	5.3	77–86 (80.95)	5.4
ASES Score	74–88 (80.45)	4.8	76–89 (81.88)	4.9
UCLA Score	24–32 (27.50)	5.6	24–32 (27.67)	5.6
VAS Score	0–3 (1.15)	0.7	0–3 (1.12)	0.8

ASES: American Shoulder and Elbow Surgeons; UCLA: University of California at Los Angeles; VAS: Visual Analog Scale

The GA group showed a significant improvement in postoperative shoulder functional outcomes over time. The Constant-Murley score was 76–85 (mean, 80.23; SD, 5.3) at 6 months and 77–86 (mean, 80.95; SD, 5.4) at 18 months, showing a slight increase during follow-up. Similarly, ASES scores increased from 74–88 (mean, 80.45; SD, 4.8) at 6 months to 76–89 (mean, 81.88; SD, 4.9) at 18 months. UCLA scores remained stable at 24–32 (mean, 27.50; SD, 5.6) at 6 months and 24–32 (mean, 27.67; SD, 5.6) at 18 months. These findings indicate that functional shoulder scores in patients operated on under GA were generally preserved or tended to improve during the follow-up period.

Patients had preoperative DM, HT, and CAD, and preoperative BMI ranged from 20 to 30 (mean, 24.1).

In Group B, 15 of the 40 patients were male and 25 were female. The youngest patient was 55 years old and the oldest was 80 years old, with a mean age of 64.28 years. Twenty-four patients had a crescent-type rotator cuff tear in the right shoulder, and 16 patients had one in the left shoulder.

At the 6-month postoperative follow-up, shoulder flexion ranged from 153 to 165 degrees (mean, 158.1), abduction from 152 to 160 degrees (mean, 157.2), internal rotation from 52 to 60 degrees (mean, 55.8), and external rotation from 56 to 63 degrees (mean, 59.8). At the 18-month postoperative follow-up, shoulder flexion ranged from 153 to 164 degrees (mean, 157.9), abduction from 151 to 158 degrees (mean, 156.5), internal rotation from 51 to 59 degrees (mean, 55.7),

and external rotation from 55 to 63 degrees (mean, 59.7), as measured by goniometry (Table 2).

Repair was performed using a single-row technique in 20 patients and a double-row technique in 20 patients. Tear size was reported as small in 6 patients, medium in 26 patients, large in 7 patients, and massive in 1 patient.

Patients' functional scores at 6 months postoperatively showed a Constant-Murley score of 79–89 (mean, 83.25), an ASES score of 77–90 (mean, 83.08), a UCLA score of 27–34 (mean, 29.38), and a VAS score of 0–2 (mean, 1.1). At 18 months postoperatively, the Constant-Murley score was 78–85 (mean, 81.30), the ASES score was 76–89 (mean, 81.80), the UCLA score was 25–33 (mean, 27.93), and the VAS score was 0–2 (mean, 1.11) (Table 4).

In the ISB group, postoperative shoulder functional outcomes were better at 6 months but declined over time to values similar to those in the GA group. These findings indicate that functional shoulder scores in patients operated on under ISB were better during the early follow-up period at 6 months but tended to decline by 18 months.

Patients had preoperative DM, HT, and CAD, and preoperative BMI ranged from 20 to 30 (mean, 24.3).

According to the results of the independent-samples Welch t-test, Constant-Murley, ASES, and UCLA scores at 6 months were significantly higher in the interscalene brachial plex-

Table 4. Postoperative shoulder function scores in patients undergoing surgery under interscalene brachial plexus block (ISB)

Functional scores and pain scales	6 th month Min-Max (Mean)	SD (6 th month)	18 th month Min-Max (Mean)	SD (18 th month)
Constant Murley Score	79–89 (83.25)	5.7	78–85 (81.30)	5.5
ASES Score	77–90 (83.08)	5.1	76–89 (81.8)	5
UCLA Score	27–34 (29.38)	5.9	25–33 (27.93)	5.5
VAS Score	0–2 (1.1)	0.8	0–2 (1.11)	0.8

ASES: American Shoulder and Elbow Surgeons; UCLA: University of California at Los Angeles; VAS: Visual Analog Scale

us block (ISB) group than in the general anesthesia (GA) group ($p<0.001$ for all) (Fig. 1). However, at 18 months, no statistically significant differences were observed between the groups in these three scores (Constant-Murley, $p=0.527$; ASES, $p=0.906$; UCLA, $p=0.528$) (Fig. 2).

A paired t-test was used for within-group comparisons.

Accordingly, in the within-group evaluation of the GA group, a significant increase was found in Constant-Murley scores between 6 and 18 months ($p<0.001$).

Similarly, a significant difference was found between the ASES scores at 6 and 18 months ($p<0.001$).

However, no statistically significant change was observed between the UCLA scores at 6 and 18 months ($p=0.051$).

When the ISB group was evaluated using the paired t-test, a statistically significant difference was observed in Constant-Murley scores between 6 and 18 months ($p<0.001$).

Similarly, a significant difference was observed between the ASES scores at 6 and 18 months ($p<0.001$).

The UCLA scores at 6 and 18 months also showed a significant difference ($p<0.001$).

The effects of time (6 and 18 months) and type of anesthesia (GA and ISB) on repeated measures were evaluated using a mixed ANOVA.

In the mixed ANOVA performed for the Constant-Murley score, the results were as follows: time, $p<0.001$; anesthesia,

$p<0.001$ (significant group difference); and time $\times$ anesthesia, $p<0.001$ (partial $\eta^2=0.557$, very strong) (Fig. 3).

In the mixed ANOVA performed for the ASES score, the results were as follows: time, $p=0.559$ (not significant); anesthesia, $p<0.001$ (significant group difference); and time $\times$ anesthesia, $p<0.001$ (partial $\eta^2=0.589$, very strong) (Fig. 4).

In the mixed ANOVA performed for the UCLA score, the results were as follows: time, $p<0.001$; anesthesia, $p=0.068$ (borderline); and time $\times$ anesthesia, $p<0.001$ (partial $\eta^2=0.640$, very strong) (Fig. 5).

All functional scores examined (Constant-Murley, ASES, and UCLA scores) were higher at 6 months; no difference was observed between the GA and ISB groups at 18 months. The interaction between time (6 and 18 months) and anesthesia was found to be statistically very strong.

Using the independent-samples t-test, no significant differences were observed between the groups (GA and ISB) in terms of age, sex, BMI, side, tear size, repair type, or anesthesia type ($p>0.05$).

When the ROM values of the GA and ISB groups were compared within each group at 6 and 18 months using a paired t-test, significant increases were observed in flexion, abduction, internal rotation, and external rotation in the GA group. In the ISB group, increases were observed in flexion and abduction, although these were more limited than those in the GA group, and significant increases were observed in




Figure 1. Distribution of Constant Murley, ASES, and UCLA scores at 6 months




Figure 2. Distribution of Constant Murley, ASES, and UCLA scores at 18 months

internal and external rotation. The changes in both groups were statistically significant ($p < 0.05$).

When ROM values were compared separately at 6 and 18 months using the independent-samples Welch t-test, flexion and abduction degrees at 6 months were higher in the ISB group than in the GA group, with statistically significant differences ($p < 0.001$ for both). Internal and external rotation degrees were also higher in the ISB group than in the GA group, with statistically significant differences ($p < 0.01$ for both).

At 18 months, all ROM degrees (flexion, abduction, internal rotation, and external rotation) were similar between the GA

and ISB groups, with no statistically significant differences ($p > 0.05$).

In the early period, at 6 months, ISB showed superiority in ROM scores, whereas in the medium term, at 18 months, no difference was observed between the ISB and GA groups.

When the VAS score was evaluated within each group using a paired t-test, it was found to have increased in both groups (GA and ISB) from 6 to 18 months, and this change was statistically significant ($p < 0.001$). As the VAS score approaches 0, patient satisfaction is thought to increase; pa-




Figure 3. Line graph showing the relationship between Constant Murley score Time (6th and 18th month) and Anesthesia type (GA and ISB)




Figure 4. Line graph showing the relationship between ASES score Time (6th and 18th month) and Anesthesia type (GA and ISB)




Figure 5. UCLA score Line graph showing the relationship between Time (6th and 18th month) and Anesthesia type (GA and ISB)

tient satisfaction was higher at 6 months postoperatively but decreased at 18 months.

When the VAS score was evaluated between the groups separately at 6 and 18 months using the independent-samples Welch t-test, the ISB group had a lower VAS score than the GA group at 6 months, and this difference was statistically significant ($p < 0.001$). At 18 months, no significant difference was observed in VAS scores between the ISB and GA groups ($p > 0.05$).

As the VAS score approaches 0, patient satisfaction is thought to increase. At 6 months postoperatively, patient satisfaction was higher in the ISB group than in the GA group, whereas at 18 months, there was no significant difference between the groups.

When a mixed ANOVA for repeated measures was performed, time (6 and 18 months) and anesthesia type (GA and ISB) were found to be significant for all ROM parameters and VAS scores ($p < 0.001$), indicating that the effect of the anesthesia technique on functional recovery was mainly limited to the early postoperative period.

Discussion

Rotator cuff tears are a common problem in orthopedic surgery, and determining the type of tear is important for treatment.

Burkhart and Lo classified rotator cuff tears into four main groups: crescent-shaped, U-shaped, L-shaped, and massive tears. Crescent-shaped tears generally have sufficient me-

dial-to-lateral mobility and can therefore be anatomically repaired directly onto the humeral tubercle under low tension.^[9] If rotator cuff tears are classified geometrically, they can be categorized as Type 1, crescent-shaped tears;^[10] Type 2, longitudinal (U- or L-shaped) tears;^[10] Type 3, massive contracted tears; and Type 4, rotator cuff arthropathy tears associated with significant glenohumeral arthrosis and complete loss of the acromioclavicular space.^[11]

Our study aimed to investigate the short- and medium-term functional changes in patients with crescent-shaped rotator cuff tears following single-row and double-row arthroscopic repairs according to the type of anesthesia.

Liu et al.^[12] evaluated the analgesic effect of a single dose of interscalene block (SISB) in the early period following arthroscopic rotator cuff repair. In patients who received SISB, pain scores on the day of surgery and especially at 6 hours were significantly lower than those in the general anesthesia group. Furthermore, insulin levels were significantly reduced in the SISB group at 42 hours postoperatively. In contrast, no significant changes were observed in DHEA-S and fibrinogen levels. The study demonstrated that SISB could be administered without complications and effectively improved early postoperative pain control.^[12]

In a study by Gurger and Ozer,^[13] the effect of the combination of general anesthesia (GA) and continuous interscalene block (CISB) on postoperative outcomes in arthroscopic rotator cuff repair was evaluated. Patients who received CISB+GA showed lower pain scores and a reduced need for additional analgesics in the early postoperative period. Additionally, Constant scores at 6 weeks were higher in the CISB+GA group. However, functional outcomes at 6 months were similar between the two groups. These findings suggest that CISB supports early pain control and functional recovery but does not provide a significant advantage in long-term outcomes.^[13]

Orhan et al.^[14] evaluated postoperative pain intensity and time to pain onset in shoulder arthroscopies performed under interscalene block. In a retrospective review of 172 patients, the type of surgery (rotator cuff repair or instability surgery) and the duration of the operation (< 30 min or ≥ 30 min) were found to have no significant effect on 24-hour VAS scores, pain onset, or the need for additional analgesics. The findings indicate that surgical and patient characteristics do not determine the postoperative pain pattern after interscalene block.^[14]

Peripheral nerve blocks (PNBs) are an important component of multimodal analgesia and can be administered as a single-dose ISB or continuous ISB. Meta-analyses of randomized controlled trials have shown that PNBs provide better pain control and lower opioid consumption than intravenous opioids in patients undergoing upper- and lower-extremity surgeries. Single-dose ISB may shorten the time to readiness for discharge compared with continuous ISB. The main limitation of single-dose ISB is that its effect is limited to 12–24 hours.^[15]

In our study, the intraoperative effects of peripheral nerve block methods and general anesthesia, postoperative vital signs, and biochemical marker monitoring were not evaluated, and only a single-dose interscalene block was applied among the peripheral nerve block methods. Evaluations were performed at the 6- and 18-month postoperative follow-ups; however, the duration of the nerve block and rebound pain effects were not examined.

Reyes-Peart et al.^[16], in a cohort study of 30 patients, showed that shoulder function improved significantly over time after rotator cuff surgery. Constant scores increased significantly from the preoperative period to the 3rd and 6th months. There was no difference in functional outcomes between patients who initially underwent arthroscopy and subsequently converted to open surgery and those who underwent arthroscopy alone. These results support the view that the surgical technique is not decisive for functional improvement and that recovery is primarily dependent on time and the rehabilitation process.^[16]

The most commonly used method today is arthroscopic repair, which was used in our study. All operations were completed arthroscopically, and patients were followed up in the short and medium term and underwent a standardized rehabilitation program.

Chen et al.^[17] conducted a meta-analysis comparing early and late rehabilitation protocols for improving shoulder function after rotator cuff repair. The study included randomized controlled trials (RCTs) selected from international and Chinese databases. According to the results of nine RCTs involving a total of 830 patients, there was no significant difference between the two approaches in terms of VAS, SST score, recovery rate, and re-tear rate at 6 and 12 months. ASES scores were similar at 6 months; the difference observed at 12 months was not clinically meaningful at the minimal clinically important difference (MCID) level. These results showed that early and late rehabilitation pro-

vided similar contributions to shoulder function. The authors concluded that when planning rehabilitation, the balance between tendon healing and range of motion should be considered, and the protocol should be flexibly adjusted according to the patient's clinical goals.^[17]

Li et al.^[18] investigated the early clinical value of enhanced recovery after surgery (ERAS) combined with an interscalene brachial plexus block (ISB) for arthroscopic rotator cuff repair (ARCR). They included 240 patients who underwent arthroscopic rotator cuff repair and randomly divided them into three groups. Groups A, B, and C were defined as surgery alone, surgery+ERAS, and ISB+surgery+ERAS, respectively. They analyzed the clinical data and postoperative indicators of the three patient groups and evaluated the early clinical effects of the ISB and ERAS protocols after ARCR. Baseline characteristics were similar between the groups. The ISB+ERAS group showed lower pain scores and shorter hospital stays during the first 24 hours after surgery. In addition, Constant-Murley and UCLA scores were higher at 6 weeks and 3 months in the groups receiving ERAS, and joint swelling was less pronounced. The incidence of postoperative stiffness did not differ between the groups. These results suggest that the ERAS protocol supports pain control and early functional recovery, while ISB provides additional benefits in pain management but does not offer additional advantages in terms of functional outcomes and complications.^[18]

In our study, all patients received standard postoperative rehabilitation consisting of passive and active shoulder movements. No differences in rehabilitation protocols were applied between the groups; therefore, no rehabilitation-related differences were observed between the groups.

Sun et al.^[19] evaluated whether acromioplasty provides additional clinical benefits in the arthroscopic repair of full-thickness rotator cuff tears in their meta-analysis. Five randomized studies involving a total of 465 patients were examined. No significant differences were found between the groups that underwent acromioplasty and those that did not in terms of ASES, Constant-Murley, and UCLA scores. Pain scores and reoperation rates were also similar between the two groups. These findings suggest that acromioplasty does not provide any additional advantage in terms of functional outcomes or pain in the short term.^[19]

In our study, acromioplasty was routinely performed arthroscopically through the lateral portal in every patient.

Gu et al.^[20] conducted a meta-analysis comparing the clinical outcomes of single-row (SR) and double-row (DR) repairs in rotator cuff injuries with different tear sizes. Data from 10 clinical studies showed that the DR technique generally provided better elevation ROM and a lower re-tear rate. There was no significant difference in clinical outcomes between SR and DR repairs for small tears (<3 cm). In contrast, for large tears (≥ 3 cm), DR repair was superior to SR repair in terms of ASES and UCLA scores. These findings suggest that the DR technique may be more advantageous in terms of functional outcomes and structural integrity, particularly for large tears.^[20]

Núñez et al.^[21] conducted a systematic review comparing the clinical outcomes of double-row and single-row techniques in arthroscopic rotator cuff repair. No demographic differences were found between the two groups in the 13 randomized studies included. The findings showed that the double-row technique was superior to the single-row technique in terms of UCLA scores. No significant differences were found in Constant-Murley and ASES scores. However, a higher tendon healing rate and a lower risk of re-tearing were observed with the double-row technique. These results support the notion that double-row repair may offer advantages in terms of structural healing and specific functional measurements.^[21]

In our study, both single-row and double-row repair techniques were used, and no significant differences were observed between the techniques in terms of ROM values, VAS, CMS, ASES, and UCLA scores.

In our study, the intraoperative effects of peripheral nerve block methods and general anesthesia, postoperative vital signs, and biochemical marker monitoring were not evaluated. Only a single-dose interscalene block was used among the peripheral nerve block methods. This method was compared with general anesthesia during postoperative follow-up in terms of functional shoulder scores, pain scales, and joint range of motion in the short and medium term (at the 6- and 18-month follow-ups).

Limitation

The limitations of our study include the small number of patients, the limited follow-up period of 18 months, and the use of only a single-dose interscalene block among peripheral nerve block anesthesia techniques. It is anticipated that a larger number of patients and a longer postoperative fol-

low-up period may reveal changes in functional scores, pain scores, and shoulder joint range of motion.

Conclusion

At 6 months postoperatively, functional shoulder scores (Constant-Murley, ASES, and UCLA scores) were better in the group that received interscalene block anesthesia than in the group that received general anesthesia, whereas at 18 months postoperatively, there was no difference between the two groups. There was no difference between the two groups in shoulder joint movements (flexion, abduction, internal rotation, and external rotation) or VAS scores at either 6 or 18 months postoperatively.

In patients who underwent ISB, the need for NSAIDs and opioid analgesics for postoperative pain management was lower, whereas it was higher in the GA group. Our study yielded results consistent with the literature, showing that peripheral nerve blocks contributed more to postoperative pain management and early rehabilitation.

Our study included only patients who underwent arthroscopic repair of crescent-shaped tears. Shoulder functional scores and range of motion were compared between general anesthesia and single-dose interscalene brachial plexus block anesthesia using different variables. It is anticipated that our results may differ when compared with those obtained using different peripheral nerve blocks, larger patient populations, different tear types and repair methods, and longer follow-up periods.

Disclosure

Ethics Committee Approval: This study was approved by The Scientific Research Ethics Committee of Balıkesir Atatürk City Hospital on 19.06.2025, with decision no. 2025/06/60.

Informed Consent: Written informed consents were obtained from patients who participated in this study.

Conflict of Interest: The authors declare that there is no conflict of interest.

Use of AI for Writing Assistance: The authors declared that artificial intelligence was not used in the study.

Financial Disclosure: The authors declared that this study received no financial support.

Authorship Contributions: Concept – A.Ç., Ö.K., B.Y.; Design – A.Ç., Ö.K., B.Y.; Supervision – A.Ç., Ö.K., B.Y.; Resource – A.Ç., Ö.K., B.Y.; Materials – A.Ç., Ö.K., B.Y.; Data Collection and/or Processing – A.Ç., Ö.K., B.Y.; Analysis and/or Interpretation – A.Ç., Ö.K., B.Y.; Literature Search – A.Ç., Ö.K., B.Y.; Writing – A.Ç., Ö.K., B.Y.; Critical Reviews – A.Ç., Ö.K., B.Y.

Peer-review: Externally peer-reviewed.

References

1. Zhang C, Wu J, Li X, Wang Z, Lu WW, Wong TM. Current biological strategies to enhance surgical treatment for rotator cuff repair. *Front Bioeng Biotechnol* 2021;9:657584.
2. Tafti D, Schultz D. Shoulder joint injection. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024.
3. Schmidt CC, Morrey BF. Management of full-thickness rotator cuff tears: appropriate use criteria. *J Shoulder Elbow Surg* 2015;24:1860–7.
4. He HB, Wang T, Wang MC, Zhu HF, Meng Y, Pan CL, et al. Tendon-to-bone healing after repairing full-thickness rotator cuff tear with a triple-loaded single-row method in young patients. *BMC Musculoskelet Disord* 2021;22:305.
5. Thangarajah T, Lo IK, Sabo MT. Rotator cuff repair techniques: Current concepts. *J Clin Orthop Trauma* 2021;17:149–56.
6. Ghodadra NS, Provencher MT, Verma NN, Wilk KE, Romeo AA. Open, mini-open, and all-arthroscopic rotator cuff repair surgery: indications and implications for rehabilitation. *J Orthop Sports Phys Ther* 2009;39:81–9.
7. Haršanji Drenjančević I, Drenjančević D, Davidović-Cvetko E, Drenjančević I, Gulam D, Kvolik S. Does the anesthesia technique affect arterial pressure and regional cerebral oxygen saturation during shoulder arthroscopy in the beach chair position? *Acta Clin Croat* 2018;57:473–9.
8. Hewson DW, Oldman M, Bedford NM. Regional anaesthesia for shoulder surgery. *BJA Educ* 2019;19:98–104.
9. Burkhart SS, Lo IK. Arthroscopic rotator cuff repair. *J Am Acad Orthop Surg* 2006;14:333–46.
10. Burkhart SS, Danaceau SM, Pearce CE Jr. Arthroscopic rotator cuff repair: Analysis of results by tear size and by repair technique-margin convergence versus direct tendon-to-bone repair. *Arthroscopy* 2001;17:905–12.
11. Davidson J, Burkhart SS. The geometric classification of rotator cuff tears: a system linking tear pattern to treatment and prognosis. *Arthroscopy* 2010;26:417–24.
12. Liu XN, Noh YM, Yang CJ, Kim JU, Chung MH, Noh KC. Effects of a single-dose interscalene block on pain and stress biomarkers in patients undergoing arthroscopic rotator cuff repair: a randomized controlled trial. *Arthroscopy* 2017;33:918–26.
13. Gurger M, Ozer AB. A comparison of continuous interscalene block versus general anesthesia alone on the functional outcomes of the patients undergoing arthroscopic rotator cuff repair. *Eur J Orthop Surg Traumatol* 2019;29:1659–66.
14. Orhan Ö, Sezgin EA, Güngör İ, Çetinkaya M, Ataoğlu MB, Kanatlı U. Interscalene block applied by an experienced anesthesiologist has a good anesthetic effect, a long duration of action, and less postoperative pain after arthroscopic shoulder procedures independent of surgery type and operation duration. *Jt Dis Relat Surg* 2023;34:445–50.
15. Joshi G, Gandhi K, Shah N, Gadsden J, Corman SL. Peripheral nerve blocks in the management of postoperative pain: challenges and opportunities. *J Clin Anesth* 2016;35:524–9.
16. Reyes-Peart R, Vargas-Vargas SF, Cruz Leal-Reyes M, Ramos-Ramos X. Functional comparison between arthroscopy and open surgery for rotator cuff injuries. *Rev Med Inst Mex Seguro Soc* 2023;61(Suppl 2):S350–5.
17. Chen Y, Meng H, Li Y, Zong H, Yu H, Liu H, et al. The effect of rehabilitation time on functional recovery after arthroscopic rotator cuff repair: a systematic review and meta-analysis. *PeerJ* 2024;12:e17395.
18. Li X, Jiang HY, Zhao YJ, Liu SZ, Pan LX. Early clinical efficacy analysis of enhanced recovery following surgery combined with interscalene brachial plexus block for arthroscopic rotator cuff repair. *Medicine (Baltimore)* 2023;102:e35943.
19. Sun Z, Fu W, Tang X, Chen G, Li J. Systematic review and Meta-analysis on acromioplasty in arthroscopic repair of full-thickness rotator cuff tears. *Acta Orthop Belg* 2018;84:54–61.
20. Gu Z, Wu S, Yang Y, Ren T, Zhang KW. Comparison of arthroscopic single-row and double-row repair for rotator cuff injuries with different tear sizes: a systematic review and meta-analysis. *Orthop J Sports Med* 2023;11:23259671231180854.
21. Núñez JH, Montenegro JD, Surroca M, Ocrospoma-Flores B, Guerra-Farfán E, Mendez-Sanchez G, et al. Arthroscopic rotator cuff repair using a single or double row technique: A meta-analysis of randomized clinical trial. *Rev Esp Cir Ortop Traumatol* 2025;69:659–69.



Assessment of Determinants Influencing Fall Risk Among Geriatric Patients

Geriatrik Hastalarda Düşme Riskini Etkileyen Faktörlerin Değerlendirilmesi

Elem Yorulmaz,¹ Berna Günay,¹ Nilgün Mesci,¹ Büşra Üstün,² Duygu Geler Külcü¹

ABSTRACT

Objectives: Falls in geriatric populations may result in significant injury and subsequent functional decline, thereby diminishing overall quality of life. Therefore, accurate assessment of fall risk and comprehensive identification of contributing factors are of fundamental importance, particularly within geriatric populations.

Methods: Geriatric patients admitted to the inpatient Physical Therapy and Rehabilitation unit were enrolled in the study. Data were collected on patients' age, sex, height, weight, educational status, medications potentially associated with falls, and assistive devices used during ambulation. Cognitive function was evaluated using the Hodkinson Abbreviated Mental Test. All participants underwent assessment with the Functional Ambulation Category (FAC), the Berg Balance Scale (BBS), and the Falls Efficacy Scale (FES). Patients were additionally questioned regarding visual impairments; urinary symptoms; nutrition-related risk factors; psychological complaints; and environmental hazards in the home that might elevate fall risk. Observations were also made to identify risky behaviors during mobility and transfers. Participants were subsequently categorized into two groups based on their fall history, and differences in these variables were analyzed between the groups.

Results: A total of 54 patients aged 65 years and older were included in the study. Patients were categorized into two groups: fallers and non-fallers. A significant difference in the female-to-male ratio was observed between the two groups ($p=0.002$). While no significant differences were observed between the two groups in terms of age, height, weight, cognitive status, FAC, or FES scores, the faller group demonstrated significantly lower BBS scores and a higher frequency of mobility- and transfer-related risky behaviors ($p=0.013$, $p=0.012$).

Conclusion: This study found that fall risk in geriatric patients was significantly associated with reduced balance and risky behaviors during mobility and transfers. The results underscore the need for systematic fall-risk assessment, close supervision, and balance-focused interventions in inpatient rehabilitation to reduce falls and support prevention strategies after discharge.

Keywords: Aged; fall risk; postural balance.

ÖZET

Amaç: Geriatrik popülasyonda görülen düşmeler, ciddi yaralanmalara ve buna bağlı fonksiyonel kayba yol açarak yaşam kalitesini belirgin biçimde azaltılabilir. Bu nedenle, düşme riskinin doğru şekilde değerlendirilmesi ve katkıda bulunan faktörlerin kapsamlı biçimde belirlenmesi, özellikle geriatrik hastalarda temel bir gerekliliktir.

Yöntem: Fizik Tedavi ve Rehabilitasyon servisine yatırılan geriatri hastaları çalışmaya dâhil edildi. Hastaların yaşı, cinsiyeti, boyu, kilosu, eğitim durumu, düşme ile ilişkili olabilecek ilaç kullanımı ve yürüme sırasında kullanılan yardımcı cihazlara ilişkin veriler kaydedildi. Kognitif durum, Hodkinson Kısaltılmış Mental Testi ile değerlendirildi. Tüm katılımcılara Fonksiyonel Ambulasyon Skalası (FAS), Berg Denge Ölçeği (BDÖ) ve Düşme Etkililiği Ölçeği (DEÖ) uygulandı. Ayrıca hastalarda görme problemleri, üriner semptomlar, beslenmeye dayalı risk faktörleri, psikolojik yakınmalar ve ev içi çevresel tehlikeler sorgulandı. Mobilite ve transfer sırasında gözlemlenen riskli davranışlar da kaydedildi. Hastalar düşme öyküsü varlığına göre iki gruba ayrıldı ve değişkenler arasındaki farklar gruplar arasında karşılaştırıldı.

¹Department of Physical Medicine and Rehabilitation, Haydarpaşa Numune Training and Research Hospital, İstanbul, Türkiye

²Department of Physical Medicine and Rehabilitation, Van Training and Research Hospital, Van, Türkiye

Cite this article as:

Yorulmaz E, Günay B, Mesci N, Üstün B, Geler Külcü D. Assessment of Determinants Influencing Fall Risk Among Geriatric Patients. Bosphorus Med J 2026;13(2):38-47.

Received: 17.11.2025

Revision: 14.01.2026

Accepted: 09.07.2026

Correspondence:

Dr. Elem Yorulmaz,
Department of Physical Medicine and Rehabilitation, Haydarpaşa Numune Training and Research Hospital, İstanbul, Türkiye

Phone:

+90 536 470 22 46

e-mail:

elem_inal@yahoo.com

OPEN ACCESS



This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License.

Bulgular: Çalışmaya 65 yaş ve üzeri toplam 54 hasta dâhil edildi. Hastalar düşen ve düşmeyen olmak üzere iki gruba ayrıldı. Cinsiyet dağılımı açısından gruplar arasında anlamlı bir fark bulundu ($p=0,002$). Yaş, boy, kilo, kognitif durum, FAS ve DEÖ skorları açısından gruplar arasında anlamlı bir fark bulunmazken, düşen grupta BDÖ skorlarının belirgin şekilde daha düşük olduğu ve mobilite ile transfer sırasında riskli davranışların daha sık görüldüğü saptandı ($p=0,013$, $p=0,012$).

Sonuç: Çalışmada, geriatri hastalarında düşme riskinin azalmış denge ile mobilite ve transfer sırasında ortaya çıkan riskli davranışlarla anlamlı düzeyde ilişkili olduğu gösterilmiştir. Bulgular, yatırılarak rehabilitasyon uygulanan hastalarda sistematik düşme risk değerlendirmesinin, yakın gözlemin ve denge odaklı müdahalelerin düşme riskinin azaltılmasında ve taburculuk sonrası korunma stratejilerinin desteklenmesinde kritik önem taşıdığını ortaya koymaktadır.

Anahtar sözcükler: Düşme riski; postural denge; yaşlı.

Falls are among the most common and preventable causes of morbidity, mortality, and loss of independence in older adults, and with increasing life expectancy worldwide, the incidence of falls has risen in the population aged 65 years and older.^[1,2] Epidemiological studies report that 25–35% of individuals over the age of 65 experience at least one fall each year, and approximately 10% of these falls result in injury.^[3] Serious consequences of falls include fractures, traumatic brain injuries, hip fractures, prolonged bed rest, functional decline, and mortality.^[4] It has been reported that one in every five falls in older adults results in a fracture or head trauma.^[1] Furthermore, fall-related mortality rates have increased dramatically over the years, and the Centers for Disease Control and Prevention (CDC) reports that in the United States, an older adult dies from a fall every 20 minutes.^[1,5] In addition to these severe health consequences, healthcare costs associated with falls are also rising globally.^[6]

An increase in fall risk is multifactorial, and individual factors such as balance impairments, visual and vestibular system problems, reduced proprioceptive sensitivity, decreased muscle strength, neuropathy, and cognitive slowing contribute to heightened susceptibility to falls.^[2,7] Among environmental factors, the home setting constitutes the most critical component. Approximately 79% of falls occur within the home environment, with bedrooms, staircases, and bathrooms reported as the most frequent locations.^[6] The WHO report emphasizes that slippery floors, inadequate lighting, cluttered household arrangements, loose rug edges, and structural inadequacies in stairways all increase fall risk.^[2] In older adults, delayed motor responses further impair the postural corrective actions required to prevent a fall, thereby increasing the likelihood of falling.^[1]

Recent systematic reviews and meta-analyses have shown that factors such as sarcopenia, mobility limitations, depression, polypharmacy, the use of antihypertensive or psychotropic medications, sleep disorders, and reduced

gait speed significantly increase fall risk.^[3,6] Sarcopenia, in particular, has been reported to both elevate fall risk and influence the severity of fall-related injuries.^[8] Weight loss, fatigue, low physical activity, slow walking speed, and reduced grip strength have also been defined as high-risk conditions for falls, disability, hospitalization, and mortality in older adults.^[9] Risky behaviors during mobility and transfers—such as rushing, failing to use necessary assistive devices, incorrect use of equipment, or wearing inappropriate footwear—have likewise been associated with an increased likelihood of falling.^[2,10]

Urinary incontinence and urgency symptoms are additional factors that heighten fall risk in older adults and have been linked to increased fall incidence in multiple studies.^[11,12]

In addition to its physical consequences, fall-related psychosocial problems are also common among older adults. Fear of falling, social isolation, and activity restriction can lead to muscle strength loss, reduced physical activity, decreased quality of life, and depression, collectively creating a vicious cycle that further increases fall risk.^[8,13] Community-based data indicate that among individuals aged 60 years and older, the prevalence of fear of falling is 69.2%, while fall-related activity restriction occurs in 38.4% of adults in this age group.^[1]

Taken together, these findings indicate that falls represent a common, multifactorial, yet preventable public health concern among older adults. Therefore, accurate assessment of fall risk, comprehensive evaluation of multiple contributing factors, and the development of individualized, evidence-based interventions are critical to reducing fall frequency and minimizing fall-related adverse outcomes.^[3] In this context, it is essential to identify the clinical, functional, psychosocial, and environmental factors that contribute to fall risk among community-dwelling older adults and to evaluate these factors within an integrated and holistic framework.^[6]

This study aims to contribute to this field by evaluating the clinical, functional, psychosocial, and environmental risk factors described in the literature among older patients hospitalized in a physical medicine and rehabilitation (PM&R) clinic and by identifying the factors that increase fall risk in this population.

Methods

This study was conducted as a single-center, cross-sectional study in the PM&R inpatient unit of a tertiary care hospital. The study protocol was approved by the Haydarpaşa Numune Training and Research Hospital Clinical Research Ethics Committee (ethics approval number: HNEAH-KAEK 2021/97, date: 15.03.2021), and all procedures were carried out in accordance with the principles of the Declaration of Helsinki. Written informed consent was obtained from all participants.

The inclusion criteria were being over 65 years of age, having received the diagnosis that led to hospitalization in the PM&R clinic at least one year prior, having a Functional Ambulation Category (FAC) level of at least 3, being able to ambulate independently despite the need for an assistive device, being able to follow commands, communicate verbally, and having no language or speech impairments that would interfere with understanding or responding to the assessment questions. Exclusion criteria included impaired speech or comprehension, the presence of limb amputation, and being classified as FAC level 1 or 2.

A total of 54 patients aged 65 years and older who were hospitalized with various diagnoses in the PM&R clinic between May 15 and August 15, 2021, were included in the study. All assessments were performed by the same clinician.

Demographic data, including age, sex, height, weight, and educational status, as well as the diagnosis leading to hospitalization and the use of assistive devices for ambulation, were recorded. In addition, medical treatments that could potentially be related to fall risk (hypnotic-sedative medications, drugs used for Parkinson's disease, diuretics, antihypertensives, and antidepressants) were reviewed, and the number of medications used from this group was documented.

Conditions known to be part of the fall etiology in the literature, as well as those considered potentially related to falls based on our clinical experience, were questioned in each patient, and responses were recorded as "present/absent."

These included visual impairments that may cause difficulty in seeing objects or navigating the environment, nutrition-related risks such as weight loss and loss of appetite, the presence of urgency, nocturia, or urinary incontinence, and environmental hazards in the home or living area that may contribute to falls, such as inappropriate rugs, inadequate lighting, or slippery floors.

In addition, the presence of depressive or anxious psychological complaints was assessed in each patient and recorded as "none/mild/moderate/severe" in terms of their impact on activities of daily living.

Patients were evaluated by the clinician for the presence or absence of risky behaviors during mobility and transfers. For the assessment of transfers, patients were asked to perform actions such as moving from the bed to a chair and from the chair back to the bed, involving coordinated movements between surfaces of different heights. Mobility was assessed by asking the patient to walk a predetermined distance within the clinic corridor. During these activities, behaviors such as rushing, not using the assistive device required for ambulation, or wearing shoes improperly were recorded as the presence of a risky behavior during the corresponding activity.

Patients' ambulation status was assessed using the FAC, which is an ordinal clinical tool used to classify ambulatory capacity on a 0–5 scale, where higher scores indicate greater independence. The scale reflects the level of assistance required during ambulation, ranging from non-functional walking (score 0) to fully independent community ambulation (score 5).^[14]

Balance was assessed using the Berg Balance Scale (BBS),^[15] a 14-item performance-based measure evaluating static and dynamic balance. Each item is scored on a 0–4 scale, yielding a total score ranging from 0 to 56, with higher scores indicating better balance performance. The Turkish version of the BBS has been culturally adapted and shown to be valid and reliable in patients with stroke.^[16]

Fear of falling was assessed using the Falls Efficacy Scale (FES), developed by Tinetti and colleagues.^[17] The FES consists of 10 items, each scored on a 1–10 scale, yielding a total score ranging from 10 to 100, with higher scores indicating greater fear of falling.

Cognitive status was assessed using the Hodkinson Abbreviated Mental Test Score (AMTS).^[18] The AMTS consists of

10 items evaluating orientation, short-term memory, and attention. Each correct answer is scored as 1 point, yielding a total score between 0 and 10, with lower scores indicating greater cognitive impairment. Patients with AMTS scores of 9–10 were considered cognitively intact, scores of 7–8 were classified as mildly impaired, scores of 5–6 as moderately impaired, and scores of 4 or below as severely impaired.

Patients were divided into two groups—fallers ($n=27$) and non-fallers ($n=27$)—based on their history of falls within the past year, and differences between the groups regarding these factors were evaluated. Demographic data, diagnoses, and functional assessment results were recorded.

Statistical Analysis

The statistical analysis of the study data was performed using SPSS Statistics version 21 software (IBM Corp., NY, USA). The normality of data distribution was assessed with the Shapiro–Wilk test. Since the data did not follow a normal distribution, descriptive statistics were presented as median (minimum–maximum) values for continuous variables and as numbers and percentages (%) for categorical variables. For comparisons between groups, the Chi-square (χ^2) test was used for categorical variables, and the Mann–Whitney U test was used for continuous variables. Correlation analysis was conducted using Spearman's rank correlation coefficient (Spearman's rho) to evaluate the relationships between variables. A p -value of <0.05 was considered statistically significant.

Results

A total of 54 patients aged over 65 years were included in the study. While half of the patients ($n=27$) had no previous history of falling, the other half had experienced at least one fall. Study data were analyzed by comparing the non-faller and faller groups.

Demographic Findings

There were no statistically significant differences between the two groups in terms of age, height, weight, or educational status ($p>0.05$). However, gender distribution differed significantly; whereas all patients in the non-faller group were women, the proportion of men in the faller group was 29.6% ($p=0.002$) (Table 1).

Clinical and Functional Assessment Findings

The comparison of clinical and functional variables between the groups is presented in Table 2. No significant differences were found between the faller and non-faller groups regarding assistive device use, presence of neurological diagnosis, medication use, psychological status, cognitive status (AMTS), FAC scores, FES scores, visual status, environmental risk factors, nutrition-related risks, or urinary incontinence ($p>0.05$). In contrast, BBS scores were significantly lower in the faller group ($p=0.003$). Risky behaviors during mobility and transfers were also observed significantly more frequently in the faller group ($p=0.013$ and $p=0.012$, respectively).

Relationships Between Variables

When the relationships between variables were examined (Table 3), a moderate and negative correlation was found between assistive device use and balance performance ($r=-0.458$, $p<0.0001$). Balance scores (BBS) showed a strong negative correlation with fear of falling (FES) ($r=-0.495$, $p<0.0001$) and a positive correlation with functional ambulation level (FAC) ($r=0.307$, $p=0.024$). BBS also demonstrated a weak but significant correlation with nutrition-related risks ($r=0.304$, $p=0.025$) and with gender ($r=-0.298$, $p=0.028$). Assistive device use was additionally correlated with FAC ($r=-0.395$, $p=0.003$) and FES ($r=0.368$, $p=0.006$).

Table 1. Demographic characteristics according to fall status

Variable	Non-fallers ($n=27$)	Fallers ($n=27$)	p
Age (years)	(62–81) (71)	(60–85) (73)	0.972*
Sex n (%)	Female: 27 (100%) Male: 0 (0%)	Female: 19 (70.4%) Male: 8 (29.6%)	0.002**
height (cm)	(146–166) (155)	(148–185) (158)	0.111*
weight (kg)	(58–100) (79)	(52–130) (80)	0.782*
Educational status n (%)	Illiterate: 3 (11.1%) Primary school: 22 (81.5%) Secondary school: 2 (7.4%)	Illiterate: 4 (14.8%) Primary school: 18 (66.7%) Secondary school: 2 (7.4%) University: 3 (11.1%)	0.315**

*:Mann–Whitney U test **: Chi-square (χ^2) test

Table 2. Clinical and functional assessments

Variable	Non-fallers n (%)	Fallers n (%)	p
Assistive device use	None: 20 (74.1%) / Single: 7 (25.9%)	None: 15 (55.6%) / Single: 9 (33.3%) / Double: 1 (3.7%) / Walker: 2 (7.4%)	0.265**
Neurological diagnosis	Yes: 2 / No: 25	Yes: 7 / No: 20	0.068**
Medication use	None: 4 (14.8%) / 1 medication: 11 (40.7%) / 2 medications: 12 (44.4%)	None: 6 (22.2%) / 1 medication: 10 (37.0%) / 2 medications: 10 (37.0%) / ≥2 medications: 1 (3.7%)	0.653**
Psychological status	None: 13 (48.1%) / Mild: 14 (51.9%)	None: 11 (40.7%) / Mild: 14 (51.9%) / Moderate: 2 (7.4%)	0.338**
Cognitive status (AMTS)	Intact: 13 (48.1%) / Mild: 10 (37.0%) / Moderate: 4 (14.8%)	Intact: 16 (59.3%) / Mild: 7 (25.9%) / Moderate: 4 (14.8%)	0.657**
FAC	(4–5) (5)	(3–5) (5)	0.123*
FES	(10–69) (31)	(6–81) (39)	0.386*
BBS	(27–51) (44)	(24–49) (39)	0.003*
Vision	Present: 4 (14.8%) / Absent: 23 (85.2%)	Present: 4 (14.8%) / Absent: 23 (85.2%)	1.000**
Mobility	Risky: 7 (25.9%) / Safe: 20 (74.1%)	Risky: 16 (59.3%) / Safe: 11 (40.7%)	0.013**
Transfers	Risky: 6 (22.2%) / Safe: 21 (77.8%)	Risky: 15 (55.6%) / Safe: 12 (44.4%)	0.012**
Environment	Risky: 3 (11.1%) / Safe: 24 (88.9%)	Risky: 1 (3.7%) / Safe: 26 (96.3%)	0.299**
Nutrition risk	No risk: 27 (100%)	Risk: 3 (11.1%) / No risk: 24 (88.9%)	0.075**
Incontinence	Present: 23 (85.2%) / Absent: 4 (14.8%)	Present: 22 (81.5%) / Absent: 5 (18.5%)	0.715**

*:Mann–Whitney U test **: Chi-square (χ^2) test, AMTS: Hodkinson Abbreviated Mental Test Score; FAC: Functional Ambulation category; FES: Falls Efficacy scale; BBS: Berg Balance scale.

A strong positive correlation was found between risky behavior during transfers and risky behavior during mobility ($r=0.696$, $p<0.0001$). Cognitive status showed a weak positive correlation with risky behavior during transfers ($r=0.303$, $p=0.026$), while the presence of a neurological diagnosis was negatively correlated with risky behavior during mobility ($r=-0.273$, $p=0.046$).

When variables associated with falls were examined, gender was positively correlated with falling ($r=0.417$, $p=0.002$). In addition, a history of falling was significantly correlated with balance scores ($r=-0.404$, $p=0.002$), risky behavior during transfers ($r=-0.342$, $p=0.011$), and risky behavior during mobility ($r=-0.337$, $p=0.013$). Gender was also negatively correlated with medication use ($r=-0.332$, $p=0.014$). Finally, there was a weak but significant negative correlation between BMI and incontinence ($r=-0.312$, $p=0.021$).

Distribution of Diagnoses

The distribution of PM&R diagnoses according to fall history is presented in Table 4. The most common diagnosis in both groups was lumbar disc herniation (LDH), observed in equal numbers in the non-faller and faller groups ($n=9$ each). Gonarthrosis was present in 4 patients in the non-faller group

and 3 patients in the faller group. Hemiplegia was more frequently observed in the faller group ($n=4$ vs. $n=1$). Impingement was identified only in the non-faller group ($n=3$), whereas brachial plexus injury and radius fracture were found exclusively in the faller group ($n=1$ each). Lumbar spondylosis was detected in five non-fallers and three fallers, while operated LDH occurred only in the faller group ($n=2$). Less common diagnoses in the non-faller and faller groups, respectively, were paraplegia ($n=1$, $n=2$), subdural hematoma (SDH) ($n=1$, $n=1$), and spinal stenosis ($n=1$, $n=1$).

Regarding the distribution of PM&R diagnoses by sex, LDH was observed in 16 women, lumbar spondylosis in 8, gonarthrosis in 5, hemiplegia in 4, and impingement in 3 patients. Additionally, paraplegia and spinal stenosis were each present in 2 women, and 1 patient each had knee arthroplasty, hip arthroplasty, brachial plexus injury, operated LDH, SDH, and radius fracture. Among men ($n=8$), the distribution was as follows: gonarthrosis ($n=2$), LDH ($n=2$), hemiplegia ($n=1$), operated LDH ($n=1$), paraplegia ($n=1$), and SDH ($n=1$).

The distribution of neurological diagnoses by sex is presented in Table 5. Neurological diagnoses were present in 15.2% of women and 25% of men.

Table 3. Correlation table (only statistically significant variables are listed)

Variable 1	Variable 2	R	p***
Assistive device use	BBS	−0.458	<0.0001
BBS	FES	−0.495	<0.0001
Risky behavior during transfers	Risky behavior during mobility	0.696	<0.0001
Sex	Falls	0.417	0.002
BBS	Falls	−0.404	0.002
Assistive device use	FAC	−0.395	0.003
Assistive device use	FES	0.368	0.006
Falls	Risky behavior during transfers	−0.342	0.011
Falls	Risky behavior during mobility	−0.337	0.013
Sex	Medication use	−0.332	0.014
BMI	Incontinence	−0.312	0.021
FAC	BBS	0.307	0.024
BBS	Nutrition risk	0.304	0.025
Cognitive status	Transfers	0.303	0.026
Sex	BBS	−0.298	0.028
Neurological diagnosis	Mobility	−0.273	0.046

***: Spearman's rank correlation test. BBS: Berg Balance scale; FES: Falls Efficacy scale; FAC: Functional Ambulation category; BMI: body mass index.

Table 4. Distribution of rehabilitation-related diagnoses according to fall history

Diagnosis	Non-fallers (n=27)	Fallers (n=27)	Total
Knee arthroplasty	1	0	1
Hip arthroplasty	1	0	1
Brachial plexus injury	0	1	1
Gonarthrosis	4	3	7
Hemiplegia	1	4	5
Impingement syndrome	3	0	3
Lumbar disc herniation	9	9	18
Lumbar spondylosis	5	3	8
Operated LDH	0	2	2
Paraplegia	1	2	3
Radius fracture	0	1	1
Subdural hematoma	1	1	2
Spinal stenosis	1	1	2
Total	27	27	54

Table 5. Distribution of neurological diagnoses by gender

Neurological diagnosis	Female (n=46)	Male (n=8)	Total
Absent	39	6	45
Present	7	2	9
Total	46	8	54

and non-fallers. The principal finding of this study is that individuals classified as fallers—those who experienced at least one fall within the past year—demonstrated significantly poorer balance, as evidenced by lower balance assessment scores. This outcome underscores the central role of postural control in fall risk. In addition, unsafe mobility patterns and high-risk behaviors during transfers were more frequently observed among fallers compared with non-fallers. Collectively, these findings imply reduced self-efficacy, impaired gait stability, and deficits in executive functioning, all of which are critical components of fall mechanisms.

Recent systematic reviews emphasize that fall risk cannot be attributed to a single determinant; rather, it reflects the interplay of multiple factors, including musculoskeletal impairments, visual–vestibular dysfunctions, medication-related effects, and environmental conditions.^[1,3,19] Nevertheless, numerous studies highlight the particularly crucial contribution of balance and gait disturbances with-

Discussion

This study examined clinical, functional, cognitive, and behavioral variables that may contribute to fall risk in geriatric patients, identifying several distinctions between fallers

in the broader spectrum of fall-related risk factors. Postural control, a core physiological function that progressively declines with advancing age, has consistently been identified as a robust predictor of fall risk.^[20] In alignment with this evidence, Colón and Emeric^[1] identified balance deficits, impaired proprioception, and mobility limitations as critical intrinsic contributors to fall susceptibility. Furthermore, meta-analytic findings indicate that balance-focused exercise interventions substantially reduce fall risk, demonstrating more pronounced effects than other exercise modalities.^[3,19,21] These findings indicate that, despite the multifactorial etiology of falls, balance impairments constitute a central target for intervention. Consistent with the prevailing evidence in the literature, the results of the present study suggest that the factors most strongly associated with fall risk are balance dysfunction, risky behaviors during mobility, and transfers.

Our study differs from most investigations of fall risk factors in geriatric populations by including a heterogeneous cohort of patients with both neurological and musculoskeletal diagnoses. We observed that patients with neurological disorders and those with a history of vertebral surgery were more likely to belong to the faller group. The increased fall risk in these patients is likely attributable to balance and gait disturbances. Consistent with our findings, the literature reports that the incidence of falls is approximately three times higher in patients with mild to moderate neurological disorders compared with those without any neurological symptoms or signs.^[22] Factors contributing to falls in this population have been reported to include behavioral determinants, such as inattention, fear of falling, and improper use of assistive devices, and environmental factors, as well as biological determinants, including impaired balance, muscle weakness, spasticity, and muscle spasms.^[23] The absence of a significant relationship between environmental factors and fall risk in our study may be attributable to the reliance on patient self-reports. Participants may have had difficulty recalling or recognizing the relevance of environmental hazards, such as slippery carpets, inadequate lighting, or wet floors.

Contrary to most reports in the literature, our study did not identify significant associations between age, female gender, or body mass index and fall risk.^[21,24–26] Conversely, male gender has been reported as a factor associated with increased fall risk. This discrepancy may be attributable to the relatively small sample size of our study, almost simi-

lar values in mean age and weight between study groups, and the proportionally higher prevalence of diagnoses potentially linked to falls within the male subgroup. Although limited, similar patterns regarding the relationship between fall risk and gender have been documented in previous research. For instance, Jehu et al.^[27] reported that men experience falls more frequently than women in their prospective cohort study, attributing this difference to factors such as reduced cognitive function, impaired judgment during movement, or slower decision-making processes in men. Additionally, Gale et al.^[28] reported that pain and balance impairments are more potent predictors of fall risk in men compared with women. The higher prevalence of male participants and balance deficits within the faller group in our study aligns with this observation. Furthermore, correlation analyses revealed that male gender was associated with polypharmacy and lower balance scores, which may help to explain the elevated fall risk observed among men in our study population.

Functional mobility assessments and gait parameters represent key determinants of fall risk in older adults. In their meta-analysis, Lusardi et al.^[29] recommended the integrated use of performance-based measures evaluating both balance and mobility to more accurately assess fall risk. Furthermore, they noted that the inclusion of self-reported indicators of depression and fear of falling alongside objective performance measures offers a more thorough and reliable method for predicting fall risk. More recently, Intaruk et al.^[30] evaluated the reliability and minimal detectable change of four commonly used functional assessments—the 10-meter walk test, the timed up-and-go test, the five-times sit-to-stand test, and the 6-minute walk test—in community-dwelling older adults at high risk of falling. They concluded that each of these assessments reflects different aspects of fall risk, providing clinicians with valuable information for evaluating and monitoring the efficacy of fall prevention interventions. Although our study did not employ these specific tests, significant correlations observed between the Functional Ambulation Scale, Berg Balance Scale scores, and the use of assistive devices further support the essential role of functional mobility in fall risk determination.

Cognitive impairment has also emerged as a significant factor associated with fall risk. It has been shown that older adults with cognitive deficits experience a higher incidence of falls compared with their cognitively intact peers, often accompanied by balance impairments and depressive symptoms.^[31,32]

Furthermore, Liu et al.'s^[33] meta-analysis highlighted that cognitive vulnerability substantially increases fall risk, particularly among individuals with diminished cognitive and physical reserves. The absence of a significant association between cognitive function and fall history in our study may be attributable to the fact that the cognitive status of the majority of participants was either intact or only mildly impaired. Nevertheless, the observed relationship between cognitive status and risky behaviors during mobility and transfers suggests that a more comprehensive and detailed assessment of cognitive function may be warranted.

Fear of falling has been recognized in the literature as an independent risk factor among older adults and has been reported to be as significant as a prior fall history in restricting activities of daily living.^[25,34] Although no direct association between Falls Efficacy Scale scores and fall history was observed in our study, the significant correlations observed between Falls Efficacy Scale scores, assistive device use, and balance performance indicate that fear of falling is closely linked to functional mobility and postural control.

Another factor associated with fall risk in geriatric patients is nutritional status. Previous studies have reported an increased risk of falls among individuals who are malnourished or at nutritional risk.^[35,36] Since the nutritional status of participants in our study was not assessed in detail, the relationship between nutrition and fall risk could not be established. Nevertheless, the observed associations between symptoms, such as loss of appetite and weight loss, and balance performance underscore the importance of comprehensive nutritional assessment in studies evaluating fall risk.

Other factors commonly associated with fall risk in geriatric populations, including polypharmacy, psychological status, visual impairments, and incontinence, were also assessed in our study; however, no significant differences were observed between fallers and non-fallers.^[1,28] This finding may be attributable to the fact that psychological status, visual impairments, and incontinence were evaluated based on patient self-reports rather than objective measures. Our limitations related to medication use are that we only investigated medications associated with falls and that we evaluated patients using more than one medication together.

Our study is valuable in that it assessed fall-related factors in a heterogeneous cohort of patients with diverse diagnoses who were receiving inpatient care in the physical therapy and rehabilitation unit of a tertiary health center. Never-

theless, it has several limitations. Although a history of falls was assessed, the frequency of falls was not recorded. The cross-sectional design makes it difficult to establish a causal relationship between variables. The relatively small sample size and single-center setting may limit the generalizability of the findings. Certain important aspects, such as detailed strength assessments and gait analysis, were not directly measured. Additionally, the exclusive reliance on self-report measures to evaluate depression and other psychosocial factors may be considered another limitation.

Future studies with larger sample sizes and multicenter designs are warranted to longitudinally monitor the incidence of falls and associated risk factors, as well as to evaluate these factors using objective measures. Such research, in conjunction with findings from intervention studies targeting specific risk factors, will generate stronger evidence to inform clinical practice and contribute to the reduction of fall-related morbidity and mortality in older adults.

Conclusion

In our study, fall risk in geriatric patients was significantly associated with balance scores, risky behaviors observed during mobility, and transfers. Conducted in an inpatient physical therapy and rehabilitation unit, this study underscores the importance of systematically assessing fall risk in patients with diverse neuromusculoskeletal diagnoses. The findings highlight the need for close monitoring during ambulation and transfers, the provision of tailored recommendations to mitigate fall risk, and the incorporation of balance and coordination exercises as integral components of the treatment plan. Such an approach not only reduces the risk of falls during hospitalization but also offers a valuable opportunity to develop strategies aimed at preventing falls in daily life after discharge.

Disclosure

Ethics Committee Approval: This study was approved by the Haydarpaşa Numune Training and Research Hospital Clinical Research Ethics Committee (Decision no: HNEAH-KAEK 2021/97, date: 15.03.2021).

Informed Consent: Informed consent was obtained from all patients.

Conflict of Interest: The authors declare that they have no conflicts of interest.

Financial Disclosure: No financial support was received for this study, and the patients received no financial compensation for their participation.

Use of AI for Writing Assistance: Artificial intelligence tools were used solely for English-language editing.

Authorship Contributions: Concept – E.Y., B.G., D.G.K.; Design – E.Y., B.G., N.M.; Supervision – D.G.K., N.M.; Resource – E.Y., B.Ü., N.M.; Materials – E.Y., B.Ü., B.G.; Data collection and/or processing – E.Y., B.G., N.M., B.Ü.; Analysis and/or interpretation – E.Y., D.G.K., B.Ü.; Literature review – E.Y., B.G., N.M.; Writing – E.Y., B.G.; Critical review – D.G.K., N.M.

Peer-review: Externally peer-reviewed.

References

- Colón-Emeric CS, McDermott CL, Lee DS, Berry SD. Risk assessment and prevention of falls in older community-dwelling adults: A review. *JAMA* 2024;331:1397–406.
- World Health Organization. WHO Global Report on Falls Prevention in Older Age. Geneva: World Health Organization; 2008.
- Pillay J, Gaudet LA, Saba S, Vandermeer B, Ashiq AR, Wingert A, et al. Falls prevention interventions for community-dwelling older adults: systematic review and meta-analysis of benefits, harms, and patient values and preferences. *Syst Rev* 2024;13:289.
- Stevens JA, Corso PS, Finkelstein EA, Miller TR. The costs of fatal and non-fatal falls among older adults. *Inj Prev* 2006;12:290–5.
- Centers for Disease Control and Prevention. Older adult falls data. Available from: <https://www.cdc.gov/falls/about/index.html>. Accessed Nov 1, 2025.
- Peel NM. Epidemiology of falls in older age. *Can J Aging* 2011;30:7–19.
- Stalenhoef PA, Diederiks JP, Knottnerus JA, Kester AD, Crebolder HF. A risk model for the prediction of recurrent falls in community-dwelling elderly: a prospective cohort study. *J Clin Epidemiol* 2002;55:1088–94.
- Yardley L, Smith H. A prospective study of the relationship between feared consequences of falling and avoidance of activity in community-living older people. *Gerontologist* 2002;42:17–23.
- Fried LP, Tangen CM, Walston J, Newman AB, Hirsch C, Gottdiener J, et al. Frailty in older adults: evidence for a phenotype. *J Gerontol A Biol Sci Med Sci* 2001;56:M146–56.
- Goh JW, Singh DKA, Mesbah N, Hanafi AAM, Azwan AF. Fall awareness behaviour and its associated factors among community dwelling older adults. *BMC Geriatr* 2021;21:226.
- Moon SJ, Kim YT, Lee TY, Moon H, Kim MJ, Kim SA, et al. The influence of an overactive bladder on falling: a study of females aged 40 and older in the community. *Int Neurourol J* 2011;15:41–7.
- Bernardes RS, Barros RDS, Silva FSD, Pequeno SA, Alves AT, Garcia PA. Urinary symptoms, falls and fear of falling in older people living with cognitive impairment. *Fisioter Mov* 2024;37:e37116.
- Lee ES, Kim B. The impact of fear of falling on health-related quality of life in community-dwelling older adults: mediating effects of depression and moderated mediation effects of physical activity. *BMC Public Health* 2024;24:2459.
- Holden MK, Gill KM, Magliozzi MR, Nathan J, Piehl-Baker L. Clinical gait assessment in the neurologically impaired. Reliability and meaningfulness. *Phys Ther* 1984;64:35–40.
- Berg KO, Wood-Dauphinee SL, Williams JI, Maki B. Measuring balance in the elderly: validation of an instrument. *Can J Public Health* 1992;83(Suppl 2):S7–11.
- Sahin F, Yilmaz F, Ozmaden A, Kotevolu N, Sahin T, Kuran B. Reliability and validity of the Turkish version of the Berg Balance Scale. *J Geriatr Phys Ther* 2008;31:32–7.
- Tinetti ME, Richman D, Powell L. Falls efficacy as a measure of fear of falling. *J Gerontol* 1990;45:P239–43.
- Hodkinson HM. Evaluation of a mental test score for assessment of mental impairment in the elderly. *Age Ageing* 1972;1:233–8.
- Guirguis-Blake JM, Perdue LA, Coppola EL, Bean SI. Interventions to prevent falls in older adults: Updated evidence report and systematic review for the US preventive services task force. *JAMA* 2024;332:58–69.
- Vaishya R, Vaish A. Falls in older adults are serious. *Indian J Orthop* 2020;54:69–74.
- Schoberer D, Breimaier HE. Meta-analysis and GRADE profiles of exercise interventions for falls prevention in long-term care facilities. *J Adv Nurs* 2020;76:121–34.
- Homann B, Plaschg A, Grundner M, Haubenhofer A, Griedl T, Ivanic G, et al. The impact of neurological disorders on the risk for falls in the community dwelling elderly: a case-controlled study. *BMJ Open* 2013;3:e003367.
- Khan A, Pujol C, Laylor M, Unic N, Pakosh M, Dawe J, et al. Falls after spinal cord injury: a systematic review and meta-analysis of incidence proportion and contributing factors. *Spinal Cord* 2019;57:526–39.
- Xu Q, Ou X, Li J. The risk of falls among the aging population: A systematic review and meta-analysis. *Front Public Health* 2022;10:902599.
- Li Y, Hou L, Zhao H, Xie R, Yi Y, Ding X. Risk factors for falls among community-dwelling older adults: A systematic review and meta-analysis. *Front Med (Lausanne)* 2023;9:1019094.
- G R Neri S, S Oliveira J, B Dario A, M Lima R, Tiedemann A. Does obesity increase the risk and severity of falls in people aged 60 years and older? a systematic review and meta-analysis of observational studies. *J Gerontol A Biol Sci Med Sci* 2020;75:952–60.
- Jehu DA, Davis JC, Barha CK, Vesely K, Cheung W, Ghag C, et al. Sex differences in subsequent falls and falls risk: A prospective cohort study in older adults. *Gerontology* 2022;68:272–9.
- Gale CR, Westbury LD, Cooper C, Dennison EM. Risk factors for incident falls in older men and women: the English longitudinal study of ageing. *BMC Geriatr* 2018;18:117.
- Lusardi MM, Fritz S, Middleton A, Allison L, Wingood M, Phillips E, et al. Determining risk of falls in community dwelling older adults: a systematic review and meta-analysis using posttest probability. *J Geriatr Phys Ther* 2017;40:1–36.
- Intaruk R, Phadungkit S, Kanpai A, Pawanta K, Srihapol N, Saengsuwan J, et al. Test-retest reliability and minimal detectable change of four functional tests in community-dwelling older adults with high risk of falls. *Turk J Phys Med Rehabil* 2024;70:164–70.
- Zhang T, Yang C, Shu G, Gao C, Ma H, Zou L, et al. The direct and mediating effects of cognitive impairment on the occurrence of falls: a cohort study based on community-dwelling old adults. *Front Med (Lausanne)* 2023;10:1190831.

32. Alkhamis BA, Elrefaey BH, Almohiza MA, Alahmari KA, Alshahrani MS, Alnakhli HH, et al. Cognition, fear, and falls: psychological predictors of balance impairment in community-dwelling older adults. *Front Psychiatry* 2025;16:1610894.
33. Liu J, Wu Y, Long Z, Zhang S, Wu S. The association between cognitive frailty and the risk of fall occurrence in older adults: a meta-analysis of cohort studies. *Front Med (Lausanne)* 2025;12:1537240.
34. Liu M, Hou T, Li Y, Sun X, Szanton SL, Clemson L, et al. Fear of falling is as important as multiple previous falls in terms of limiting daily activities: a longitudinal study. *BMC Geriatr* 2021;21:350.
35. Eckert CD, Tarleton EK, Pellerin J, Mooney N, Gell NM. Nutrition risk is associated with falls risk in an observational study of community-dwelling, rural, older adults. *J Aging Health* 2022;34:1125–34.
36. Mziray M, Nowosad K, Śliwińska A, Chwesiuk M, Małgorzewicz S. Malnutrition and fall risk in older adults: a comprehensive assessment across different living situations. *Nutrients* 2024;16:3694.



Performance of the 'Turkish Triage System' During the Covid-19 Pandemic

COVID-19 Pandemisinde 'Türk Triaj Sistemi'nin Performansı

● Selman Faruk Akyıldız, ● Fatma Sarı Doğan, ● Tuba Cimilli Öztürk,
● Ebru Ünal Akoğlu, ● Yasin Bostancı

ABSTRACT

Objectives: In Türkiye, a triage system based on a 3-color code (red: very urgent, yellow: urgent, green: non-urgent) is mandatory in all public hospitals. This study aimed to analyze the triage codes assigned to confirmed COVID-19 patients according to their outcomes to test the reliability of the triage system. Our study provides data on our triage system, whose reliability has not been tested before.

Methods: Patients with positive PCR tests who were admitted to the emergency department between March 15 and December 31, 2020, were included in the study. Assigned triage codes and outcomes were analyzed retrospectively.

Results: Of the 7040 patients, 4176 (59.3%) were coded as 'green' and 333 (4.7%) were coded as 'red' at triage. The sensitivity of green triage for recovery was 60.29%, and the specificity was 78.78%. The sensitivity of red triage for critical outcomes and mortality was 57.14%, and the specificity was 96.59%. Being over 58 years old, regardless of gender, was a predictive factor for critical outcomes and mortality.

Conclusion: Based on this study, it is possible to state that the 'Turkish Triage System' is successful in recognizing patients in critical situations, but its performance in distinguishing cases with recovery is moderate in the COVID-19 patient group.

Keywords: COVID-19; emergency department; patient outcomes; triage; Turkish triage system.

ÖZET

Amaç: Türkiye'de, 3 renkli kodlamaya dayalı bir triaj sistemi (kırmızı: çok acil, sarı: acil, yeşil: acil değil) tüm kamu hastanelerinde zorunlu olarak kullanılmaktadır. Bu çalışma, triaj sisteminin güvenilirliğini test etmek amacıyla doğrulanmış COVID-19 hastalarına verilen triaj kodlarını hasta sonuçlarına göre analiz etmeyi hedeflemektedir. Çalışmamız, daha önce güvenilirliği değerlendirilmemiş olan triaj sistemimize ilişkin veriler sunmaktadır.

Yöntem: 15 Mart–31 Aralık 2020 tarihleri arasında acil servise başvuran ve PCR testi pozitif olan hastalar çalışmaya dâhil edildi. Verilen triaj kodları ve hasta sonuçları retrospektif olarak analiz edildi.

Bulgular: Toplam 7040 hastanın 4176'sı (%59,3) triajda "yeşil", 333'ü (%4,7) ise "kırmızı" olarak kodlandı. İyileşmeyi öngörmede yeşil triajın duyarlılığı %60,29, özgüllüğü ise %78,78 idi. Kritik sonuçları ve mortaliteyi öngörmede kırmızı triajın duyarlılığı %57,14, özgüllüğü ise %96,59 olarak bulundu. Cinsiyetten bağımsız olarak 58 yaşın üzerinde olmak, kritik sonuçlar ve mortalite için prediktif bir faktör olarak saptandı.

Sonuç: Bu çalışma ile "Türk Triaj Sistemi"nin kritik durumdaki hastaları tanımda başarılı olduğu; ancak COVID-19 hasta grubunda iyileşmeyle sonuçlanan olguları ayırt etmedeki performansının orta düzeyde olduğu söylenebilir.

Anahtar sözcükler: Acil Servis; COVID-19; hasta sonuçları; triaj; Türk triaj sistemi.

*This study was a thesis for graduation from Emergency Medicine Residency Program in University of Health Sciences Fatih Sultan Mehmet Training and Research Hospital.

Department of Emergency Medicine, University of Health Sciences Fatih Sultan Mehmet Training and Research Hospital, İstanbul, Türkiye

Cite this article as:

Akyıldız SF, Sarı Doğan F, Cimilli Öztürk T, Ünal Akoğlu E, Bostancı Y. Performance of the 'Turkish Triage System' During the Covid-19 Pandemic. Bosphorus Med J 2026;13(2):48-55.

Received: 18.12.2025

Revision: 14.04.2026

Accepted: 29.04.2026

Correspondence:

Dr. Selman Faruk Akyıldız,
Department of Emergency Medicine, University of Health Sciences Fatih Sultan Mehmet Training and Research Hospital, İstanbul, Türkiye

Phone:

+90 543 625 37 47

e-mail:

sfarukakyildiz@gmail.com

OPEN ACCESS



This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License.

In emergency departments (EDs), different triage systems are used in different countries to determine the urgency of patients' conditions and their priority for examination. The Emergency Severity Index (ESI), Canadian Triage System, Manchester Triage System (MTS), and Australian Triage System (ATS) are examples of triage systems used worldwide.^[1,2] Most of these systems determine the triage category using parameters such as the patient's symptoms and vital signs (blood pressure, temperature, heart rate, oxygen saturation, and respiratory rate [RR]) and the resources required to manage the patient.

In our country, triage practice became mandatory in EDs with the publication of the 'Communiqué on Implementation Procedures and Principles of Emergency Departments in Inpatient Health Facilities' by the Ministry of Health of the Turkish Government in 2009. In the specified triage system, patients are classified using three colour codes: red (very urgent), yellow (urgent), and green (less urgent), and each colour contains two subcategories.^[3] This colour-coded triage system applied in our country is similar to the MTS in terms of the urgency represented by the colours and the ATS in terms of its categories.^[4-6] The triage system that we use has not yet been named, so we will call it the 'Turkish Triage System' (TTS). We know of no published study on the reliability and validity of the triage system used in our country before or during the pandemic. Therefore, our study presents new data.

Due to the COVID-19 pandemic, EDs had to be physically reorganized to manage the burden of cases and isolate patients at risk of infection. The infection prevention and control recommendations of the World Health Organization emphasized that effective triage and isolation were needed to prevent unnecessary exposure of those at risk of contracting COVID-19.^[7] In Türkiye, a guideline containing recommendations from the Scientific Advisory Board of our Ministry of Health is used for the management and treatment of COVID-19 patients, and it has been updated periodically with new scientific evidence. Unfortunately, while there was agreement on the isolation procedures, no specific system or procedure for the triage of patients was defined, even in the latest updates.^[8,9]

During the pandemic, we applied a two-step triage system in the ED of our hospital. Suspected COVID-19 patients (based on respiratory symptoms, fever, and close contact with a person with confirmed COVID-19) were separated from other patients and later evaluated in an isolated area of the

ED. Patients in both groups were then subjected to secondary triage and classified according to the standard colours (green, yellow, or red) of the TTS.

In this study, we conducted a retrospective analysis of the triage codes of patients with confirmed COVID-19 to evaluate the performance of the TTS in distinguishing COVID-19 patients at high risk of death from those who could be treated as outpatients. We investigated disease-related mortality, intubation, intensive care hospitalization within 30 days, and recovery from the disease. Additionally, the relationship between vital signs at admission and patient outcomes was evaluated.

Methods

Study Design

We conducted a retrospective cross-sectional data analysis. Triage is conducted by two experienced nurses in the ED. One obtains information about the patient's symptoms and medical history and enters the data electronically, while the other measures the patient's vital signs. Triage nurses are selected from among nurses with at least two years of ED experience and are trained annually using simulation scenarios.

During the pandemic, patients with possible COVID-19 were isolated from other patients, and secondary triage was applied to both groups. The Glasgow Coma Score (GCS), blood pressure (BP), pulse, oxygen saturation, respiratory rate (RR), and temperature of each patient with possible COVID-19 were recorded. Patients with possible COVID-19 who had a history of COVID-19 contact and/or symptoms highly suggestive of COVID-19 infection, such as cough, shortness of breath, and fever, and who had stable vital signs were classified at the 'green' triage level and evaluated in the pandemic outpatient clinic. The remaining patients were coded as 'yellow' or 'red' triage levels according to the triage system criteria and were examined and treated separately in the appropriate areas.^[5] The triage forms recorded in the hospital's automated system were examined retrospectively.

Settings and Participants

Our hospital is a training and research hospital located in Istanbul, the city with the highest number of COVID-19 cases in Türkiye. The hospital has a tertiary ED with an average of 270,000 patient admissions annually before the pandemic. During the pandemic, the number of emergency department admissions sometimes increased three- or four-fold.

Patients who were admitted to the ED of Fatih Sultan Mehmet Training and Research Hospital between 15 March and 31 December 2020 were identified through the General Directorate of Public Health Laboratory Information Management System, in which nationwide COVID-19 polymerase chain reaction (PCR) results were recorded.

COVID-19 PCR-positive patients were evaluated retrospectively in the study. Patients who visited the ED for screening tests only and were not admitted through the emergency triage department, as well as those who underwent repeated tests during hospitalization, were excluded from the study.

Outcome Measurements

Intubation and intensive care admission were considered critical outcomes. Triage systems are primarily designed to identify or safely exclude patients with critical illness rather than to predict recovery. At times when there are excessive numbers of patient admissions, it is important to identify patients who will not require immediate ED evaluation, as well as critically ill patients and those at risk of mortality. Therefore, we focused on red and green codes. Patient outcomes were also obtained from digital records. Each patient was counted once, even if the patient died after being intubated and hospitalized in the intensive care unit because of COVID-19 within 30 days. Information that was not available in hospital records was obtained from the patients or their relatives by phone.

Mortality information was obtained using the Death Notification System or by phone. The relationship between the triage codes assigned to patients and the outcomes was statistically analysed.

Statistical Analysis

IBM SPSS Statistics 22 (IBM SPSS, Türkiye) and MedCalc version 23.5.2 (MedCalc Software Ltd) were used for statistical analysis. The Shapiro–Wilk test indicated that the data did not conform to a normal distribution ($p < 0.05$). In addition to descriptive statistical methods (mean, standard deviation, and frequency), the Mann–Whitney U test was used to compare quantitative data between the two groups. The chi-square test was used to compare qualitative data. We considered $p < 0.05$ to indicate statistical significance. In addition, specificity and sensitivity analyses were conducted using cross-tabulation. Receiver operating characteristic (ROC) curve analyses were performed to determine the efficacy of age and vital signs in predicting critical outcomes and mortality.

This study was conducted in accordance with the principles of the Declaration of Helsinki. The study was approved by the Istanbul Fatih Sultan Mehmet Training and Research Hospital Clinical Research Ethics Committee (Date: 11.06.2020, Decision no: 2020/11). Owing to the retrospective design of the study, the requirement for informed consent was waived by the Ethics Committee.

Results

A total of 7040 patients with positive COVID-19 PCR tests were included in the study. Of them, 3784 (53.8%) were male and 3256 (46.2%) were female. The patients' ages ranged from 1 to 99 years (median age, 41 years), with a mean age of 41.94 ± 15.28 years. Figure 1 depicts a flowchart of the study.

There was a statistically significant difference between the mean ages of patients with different outcomes; the mean




Figure 1. Flow chart of the study.

Table 1. ROC Curve Analyses of Vital Parameters

Parameter	Area Under Curve (%95 CI)	Cut-off Value	Sensitivity	Specificity	p
Age (year)	0.922 (0.916–0.929)	>58	82.9	87.7	<0.0001
Temperature (°C)	0.642 (0.631–0.653)	>36,6	50.3	72.7	<0.0001
Heart Rate (beats/min)	0.540 (0.528–0.552)	>88	68.6	40.9	0.067
SBP (mmHg)	0.634 (0.622–0.645)	>135	44	81.5	<0.0001
DBP (mmHg)	0.504 (0.492–0.516)	≤60	15.4	91.7	0.8651
Oxygen Saturation (%)	0.824 (0.815–0.833)	≤96	74.2	76.1	<0.0001
Respiratory Rate (breaths/min)	0.626 (0.615–0.638)	>20	28.6	99.1	<0.0001

age of patients who died (73.61±13.12 years) was significantly higher than that of those who were discharged and recovered (41.36±14.69 years) ($p<0.001$). No statistically significant effect of gender on outcome was found ($p=0.886$).

We also investigated vital signs at the time of triage. Fever, systolic blood pressure (SBP), and RR were significantly higher in patients with critical outcomes and mortality than in those who were discharged and recovered ($p<0.001$), whereas GCS and oxygen saturation were significantly lower ($p<0.001$). The cut-off values calculated for age and oxygen saturation were statistically significant. However, the cut-

off determined for oxygen saturation, ≤96%, was not considered clinically significant. The cut-off for age, >58 years, might be both clinically and statistically significant. Table 1 shows the results of the ROC curve analyses performed to determine the efficacy of age and vital signs in predicting critical outcomes and mortality.

Of the 7040 patients, 4176 (59.3%) were coded as 'green' and 333 (4.7%) were coded as 'red' at the time of triage. Patients were evaluated according to their final outcomes. The distributions of the patients and the final outcomes evaluated are summarized in Table 2. Table 3 shows comparisons of outcomes according to triage codes.

Table 2. Distribution of the patients and final outcomes

	n	%
Triage		
Green	4176	59.3
Yellow	2531	36.0
Red	333	4.7
Ward admission		
Yes	798	11.3
No	6242	88.7
ICU admission		
Yes	152	2.2
No	6888	97.8
Intubation		
Yes	100	1.4
No	6940	98.6
Exitus		
Yes	127	1.8
No	6913	98.2
Outcome		
Critical outcome and mortality	175	2.4
Recovery	6865	97.5
Total	7040	100.0

The sensitivity of the green triage code for recovery was 60.29% (95% CI: 59.12–61.45%), and the specificity was 78.78% (95% CI: 72.05–84.66%). Importantly, the negative predictive value of the green triage code for 'critical outcomes and mortality' was 0.95 (95% CI: 0.947–0.954), indicating that the green code was effective in safely excluding critical illness. The positive predictive value of the green triage code for recovery was 0.975 (95% CI: 0.971–0.978), and the positive likelihood ratio was 2.85 (95% CI: 2.14–3.80). The negative likelihood ratio was 1.99 (95% CI: 1.83–2.16).

Table 3. Comparison of the outcome against triage

		Outcome		Total	p
Triage Code		Critical outcome and mortality (n=175)	Recovery (n=6865)		
Green	n (%)	37 (0.8)	4139 (99.1)	4176 (100)	<0.0001
Yellow	n (%)	39 (1.5)	2492 (98.4)	2531 (100)	
Red	n (%)	99 (29.7)	234 (70.2)	333 (100)	
Total	n (%)	175 (2.4)	6865 (97.5)	7040 (100)	

The sensitivity of the red triage code for 'critical outcomes and mortality' was 56.57% (95% CI: 48.88–64.03%), and the specificity was 96.59% (95% CI: 96.14–97.01%). The positive likelihood ratio was 16.60 (95% CI: 13.85–19.89).

The sensitivity of the red triage code for 'critical outcomes and mortality' in the ED was 92.00% (95% CI: 73.97–99.02%), and the specificity was 95.57% (95% CI: 95.06–96.04%). The positive likelihood ratio was 20.75 (95% CI: 17.71–24.32) (Table 1).

Discussion

The increase in the number of COVID-19 cases placed a significant strain on healthcare systems worldwide. One of the most important issues was the management of COVID-19 patients who were admitted to the ED. Experts from many different healthcare centres have shared their experiences with ED organization.^[10–12] However, to our knowledge, no published studies have validated the performance of various pre-pandemic ED triage systems used during the pandemic. Additionally, the TTS had not been validated, even before the pandemic, although it has been used nationwide since 2009. We compared our results with those of performance studies of other triage systems conducted before the pandemic. Therefore, our study presents new data.

As a result of the healthcare system implemented in our country, daily ED admissions are approximately 1000 patients in almost every hospital, and this number occasionally reaches 1500–2000 in many hospitals. In Türkiye, patients who are not experiencing an emergency can easily be admitted to any level of ED, even in university hospitals, without any outpatient qualification. There is no legal obstacle to this; in fact, it is forbidden to refuse these patients. We are very uncomfortable with this situation because it is generally abused. It is ironic that we, as emergency physicians in Türkiye, had the advantage of working under such a burden as part of our daily routine during the pandemic. We think that the TTS is a beneficial tool for managing crowds in EDs.

Of the patients included in our study, 4176 (59.3%) were assigned green triage codes, 2531 (36%) yellow, and 333 (4.7%) red. Wuerz et al.^[2] evaluated the accuracy and reliability of the ESI triage system; patients were classified as very urgent (13%), urgent (52%), or not urgent (34%). They reported that this scoring was reliable in estimating the length of hospital stay compared with hospitalization, resource use, and tri-

age scoring. Zachariasse et al.^[13] studied 288,633 patients at three hospitals for one year and evaluated the validity of the MTS. The sensitivity of the MTS was 0.47–0.87 among the three hospitals, and the specificity was 0.84–0.94; the validity of the system was moderate to good. In our study, the sensitivity of the green triage code for recovery was 60.29%, and the specificity was 78.78%. Notably, the negative predictive value of the green code for critical outcomes and mortality was 0.95, suggesting that the green code was effective in safely ruling out critical illness. The sensitivity of red triage for critical outcomes and mortality was 56.57%, and the specificity was 96.59%; the positive likelihood ratio was 16.60. When we calculated the sensitivity of red triage for critical outcomes and mortality in the ED, the sensitivity increased to 92%, and the specificity was 95.57%.

In comparison with related studies, the triage system used in our hospital performed well in identifying critical cases. However, it is important to note that triage systems are primarily designed to identify or safely exclude patients with critical illness rather than to predict recovery. The moderate sensitivity of the green code for recovery is therefore expected and does not diminish the clinical utility of the system.^[2,13] This is probably due to the nature of the specific disease. Nearly half of the patients were assigned yellow triage codes. These results suggest that triage nurses may also have been reluctant to assign the green triage code because of uncertainty about the disease. We suggest that a study conducted during a non-pandemic period would be more demonstrative and realistic.

Our study, with its large number of patients, also contributes epidemiological data regarding the COVID-19 pandemic. The mean age of the patients included in our study was 41.94 ± 15.28 years, and 53.8% were male. İlhan Alp et al.^[14] studied patients with a mean age of 40.4 ± 15.67 years, similar to our study, and 57.4% of the participants were male. Ankaralı et al.^[15] reported a significant relationship between age and death due to COVID-19: the death rate increased significantly with increasing age. Ek et al.^[16] investigated the effects of COVID-19 on older people, and their sample included 85 countries with more than 5000 cases. They reported that the older population constituted a significant proportion of deaths due to COVID-19. Our study found that the mean age of patients who died (73.61 ± 13.12 years) was significantly higher than that of those who were discharged and recovered (41.36 ± 14.69 years), which was consistent with findings reported in the literature. Fifty-eight years

of age was determined to be a cut-off value for critical outcomes and mortality.

In this study, no statistically significant difference in outcomes was found between genders. The rates of 'critical outcomes and mortality' and recovery were similar in patients of both genders. Although some studies have reported a gender difference in terms of mortality and outcomes in patients with COVID-19, other studies have reported results similar to those of our study. Mohamed et al.^[17] found that death rates due to COVID-19 were similar in men and women and that there was no significant difference between the groups. However, in a study conducted in Türkiye, 66.1% of those who died were men and 33.9% were women, and the incidence of cases was 9% higher in men than in women.^[18]

In our study, temperature, systolic BP, and RR were significantly higher in COVID-19 patients whose outcomes were critical and/or fatal than in those who were discharged and recovered, whereas GCS and oxygen saturation were significantly lower. Choron et al.^[19] reported that fever was one of the parameters that could be used to estimate mortality in mechanically ventilated COVID-19 patients. Caillon et al.^[20] reported that high systolic BP at the time of hospital admission was an important risk factor in models predicting the outcomes of COVID-19 patients. Knight et al.^[21] reported that a high RR was an important parameter in predicting COVID-19-related deaths.

Of the patients evaluated in our study, 2.2% ($n=152$) were admitted to the intensive care unit, and 31.6% of them were discharged and recovered. Korkmaz reported that 10% of COVID-19 patients were admitted to the intensive care unit.^[22] Of the 1.3 million patients with COVID-19 reported to the United States Centers for Disease Control and Prevention by the end of May 2020, 2% were hospitalized in intensive care, and 5% died.^[23] Kim et al.^[24] studied 2491 COVID-19 patients and reported that the rate of hospitalization in the intensive care unit was 32%, and the rate of requiring mechanical ventilation was 19%. During the pandemic, many countries attempted to intervene using the resources available for patient care. In addition, mean age, gender distributions, and population densities vary among countries. Such factors may have affected intensive care hospitalization rates. We found that 1.4% ($n=100$) of patients were intubated, of whom 92 (92%) died and 8 (8%) were discharged and recovered. The weekly COVID-19 status reports published by our Ministry of Health reported that 8.8% of patients were intubated as of October 4, 2020, while the intubation rate between Octo-

ber 19 and October 25, 2020, was 3.8%.^[25,26] Ertürk reported that the intubation rate was 8.9%.^[27] Intubation rates in our study were lower than those reported in the literature. We suggest that one reason for this was the establishment of an 'ED critical care unit' in our hospital during the pandemic; in this unit, non-invasive mechanical ventilation was used. Additionally, patients who might otherwise have required intubation because of transfers to distant hospitals could be treated here without being intubated. Another reason might have been the advocacy of early intubation at the beginning of the pandemic. Since our study was conducted toward the middle of the pandemic, intubation rates may have been lower during this period.

Limitations

Our study is a retrospective cross-sectional study conducted at a single centre. Although single-centre studies provide detailed institutional data, they may limit the generalizability of the findings to other settings with different patient populations, triage practices, and healthcare infrastructure. Additionally, although triage is performed primarily by experienced ED nurses trained in triage, differences in their levels of experience and interpersonal dynamics may potentially affect the accuracy and consistency of triage decisions. Unfortunately, inter-rater and intra-rater agreement analyses could not be performed because of the retrospective nature of the study and the pandemic.

It is important to consider contextual factors when interpreting the results. Changes in healthcare policies and guidelines, as well as fluctuations in the number and severity of COVID-19 cases, can affect the number of patient admissions and the clinical profiles of patients seen in EDs. Similarly, the number of patients and their clinical profiles may have varied according to the virus variant circulating in the population at a given time.

In our study, only the triage of patients with PCR positivity on the relevant date was examined. However, it is now known that there are many cases in which PCR positivity is not detected at admission. Therefore, it is possible that, during the study period, many more COVID-19 patients were admitted to the ED but were excluded from the study because they were PCR-negative.

Patients with clinical findings of COVID-19 but negative PCR tests and patients with negative PCR tests but typical COVID-19 findings on thoracic CT were not included in this study.

Conclusion

Our study suggests that the TTS is successful in identifying critical outcomes and mortality in patients with COVID-19 and in safely excluding critical illness among patients assigned the green triage code. As triage systems are primarily designed to identify or safely exclude critical illness rather than to predict recovery, the moderate sensitivity of the green code for recovery does not diminish the clinical value of the system.

Our study provides valuable insight into the system's ability to identify critical illness and predict outcomes in COVID-19 patients. Further validation studies are needed after the pandemic to confirm the system's usefulness in the triage of general ED patients.

Although it has not yet been scientifically proven, based on our clinical experience, we believe that this triage system, which we have been using safely in EDs since 2009, can be a useful tool for managing patient loads during extraordinarily busy periods in EDs beyond the COVID-19 pandemic.

Disclosures

Ethics Committee Approval: This study was approved by the Istanbul Fatih Sultan Mehmet Training and Research Hospital Clinical Research Ethics Committee (Date: 11.06.2020, Decision no: 2020/11).

Informed Consent: Owing to the retrospective design of the study, the requirement for informed consent was waived by the Ethics Committee.

Conflict of Interest: None declared.

Financial Disclosure: The author declared that this study has received no financial support.

Use of AI for Writing Assistance: None declared.

Authorship Contributions: Concept – T.C.Ö.; Design – T.C.Ö.; Supervision – F.S.D.; Resource – S.F.A.; Materials – E.Ü.A.; Data collection and/or processing – S.F.A.; Analysis and/or interpretation – Y.B.; Literature review – S.F.A.; Writing – S.F.A.; Critical review – F.S.D.

Peer-review: Externally peer-reviewed.

References

- Beveridge R, Ducharme J, Janes L, Beaulieu S, Walter S. Reliability of the Canadian emergency department triage and acuity scale: interrater agreement. *Ann Emerg Med* 1999;34:155–9.
- Wuerz RC, Milne LW, Eitel DR, Travers D, Gilboy N. Reliability and validity of a new five-level triage instrument. *Acad Emerg Med* 2000;7:236–42.
- Yataklı sağlık tesislerinde acil servis hizmetlerinin uygulama usul ve esasları hakkında tebliğ. T.C. Resmi Gazete. 20 Şubat 2018-Sayı: 30338. Available at: <https://www.resmigazete.gov.tr/eskiler/2022/09/20220913-5.htm> Accessed March 27, 2021[in Turkish]
- Fernandes CM, Tanabe P, Gilboy N, Johnson LA, McNair RS, Rosenau AM, et al. Five-level triage: a report from the ACEP/ENA Five-level Triage Task Force. *J Emerg Nurs* 2005;31:39–50.
- Gilboy N, Tanabe P, Travers D, Rosenau AM. *Emergency Severity Index (ESI): A Triage Tool for Emergency Department Care. Version 4. Implementation Handbook*. Rockville (MD): Agency for Healthcare Research and Quality; 2012. AHRQ Publication No. 12-0014.
- Şimşek DÖ. Triyaj sistemlerine genel bakış ve Türkiye’de acil servis başvurularını etkileyen faktörlerin lojistik regresyon ile belirlenmesi. *Sosyal Güven* 2018;84–115.
- World Health Organization. Clinical management of severe acute respiratory infection when Novel coronavirus (2019-nCoV) infection is suspected: interim Guidance. Available at: <https://iris.who.int/items/329bc02b-d963-4e6c-a1b4-625a65b217b6> Accessed 28 Jan 2020.
- T.C. Sağlık Bakanlığı. COVID-19 (SARS-CoV-2 enfeksiyonu) rehberi. Available at: <https://acilafet.saglik.gov.tr/Eklenti/37175/0/covid-19rehberipdf.pdf> Accessed 27 March 2021 [in Turkish]
- T.C. Sağlık Bakanlığı. COVID-9 vaka sorgulama kılavuzu. Available at: https://ohsad.org/wp-content/uploads/2020/04/COVID19-VAKA-SORGULAMA-KILAVUZU-A4_1.pdf Accessed 28 Jan, 2021. [in Turkish]
- Nadarajan GD, Omar E, Abella BS, Hoe PS, Do Shin S, Ma MH, et al. A conceptual framework for Emergency department design in a pandemic. *Scand J Trauma Resusc Emerg Med* 2020;28:118.
- Schreyer KE, Del Portal DA, King LJJ, Blome A, DeAngelis M, Stauffer K, et al. Emergency Department Management of the Covid-19 Pandemic. *J Emerg Med* 2020;59:946–51.
- Af Ugglas B, Skyttberg N, Wladis A, Djärv T, Holzmänn MJ. Emergency department crowding and hospital transformation during COVID-19, a retrospective, descriptive study of a university hospital in Stockholm, Sweden. *Scand J Trauma Resusc Emerg Med* 2020;28:107.
- Zachariasse JM, Seiger N, Rood PP, Alves CF, Freitas P, Smit FJ, et al. Validity of the Manchester Triage System in emergency care: A prospective observational study. *PLoS One* 2017;12:e0170811.
- İlhan Alp S, Khayri U, Demirkıran A. Coronavirüs hastalığı 2019 (COVID-19) olgularının klinik özellikleri ve obstruktif uyku apne sendromu ile ilişkisi: kesitsel bir çalışma. *Abant Med J* 2021;10:247–57. [Article in Turkish]
- Ankaralı H, Ankaralı S, Erarslan N. COVID-19, SARS-CoV2, enfeksiyonu: güncel epidemiyolojik analiz ve hastalık seyrinin modellenmesi. *Anadolu Klin* 2020;25:1–22. [Article in Turkish]
- Ek S, İlhanlı H, Kahraman SÖ. COVID-19’un zayıf halkası: Yaşlı nüfus. *Türk Coğrafya Dergisi* 2020;(76):33–44. [Article in Turkish]
- Mohamed MO, Gale CP, Kontopantelis E, Doran T, de Belder M, Asaria M, et al. Sex differences in mortality rates and underlying conditions for covid-19 deaths in england and wales. *Mayo Clin Proc* 2020;95:2110–24.
- Yıldırım, C. COVID-19: cinsiyet bağlamında değerlendirme. *Sosyal Hizmet* 2020;94–7. [Article in Turkish]

19. Choron RL, Butts CA, Bargoud C, Krumrei NJ, Teichman AL, Schroeder ME, et al. Fever in the ICU: a predictor of mortality in mechanically ventilated COVID-19 patients. *J Intensive Care Med* 2021;36:484–93.
20. Caillon A, Zhao K, Klein KO, Greenwood CMT, Lu Z, Paradis P, et al. High systolic blood pressure at hospital admission is an important risk factor in models predicting outcome of covid-19 patients. *Am J Hypertens* 2021;34:282–90.
21. Knight SR, Ho A, Pius R, Buchan I, Carson G, Drake TM, et al. Risk stratification of patients admitted to hospital with covid-19 using the ISARIC WHO Clinical Characterisation Protocol: development and validation of the 4C Mortality Score. *BMJ* 2020;370:m3339.
22. Korkmaz N, Karakaya Z, Acar H, Bilgin S, Kayali A, et al. Acil serviste COVID-19 hastalarının prognoz tahmininde erken uyarı skorlamalarının kullanımı. *Journal of Contemporary Medicine* 2023;13:90–5. [in Turkish]
23. Stokes EK, Zambrano LD, Anderson KN, Marder EP, Raz KM, El Burai Felix S, et al. Coronavirus Disease 2019 Case Surveillance - United States, January 22-May 30, 2020. *MMWR Morb Mortal Wkly Rep* 2020;69:759–65.
24. Kim L, Garg S, O'Halloran A, Whitaker M, Pham H, Anderson EJ, et al. Risk factors for intensive care unit admission and in-hospital mortality among hospitalized adults identified through the US coronavirus disease 2019 (COVID-19)-associated hospitalization surveillance network (COVID-NET). *Clin Infect Dis* 2021;72:e206–e14.
25. T.C. Sağlık Bakanlığı. COVID-19 haftalık durum raporu:28/09/2020–/10/2020. Available at: https://covid19.saglik.gov.tr/Eklenti/39039/0/covid-19-haftalik-durum-raporu---40-haftapdf.pdf?_tag1=5DA7702AE03EEB2E875C667B8755DDDC03FD259D Accessed 3 March 2021. [in Turkish]
26. T.C. Sağlık Bakanlığı. COVID-19 Haftalık Durum Raporu:19/10/2020 – 25/10/2020. [in Turkish]
27. Ertürk NC, Turan MO. COVID-19 hastalarında beden kitle indeksinin prognoza olan etkisi. *Uzmanlık Tezi. İzmir: İzmir Kâtip Çelebi Üniversitesi Tıp Fakültesi, Atatürk Eğitim ve Araştırma Hastanesi; 2021.*



Comparison of the Efficacy of Low-Level Laser Therapy and Ultrasound Treatment in Patients with Shoulder Pain

Omuz Ağrılı Hastalarda Düşük Yoğunluklu Lazer ve Ultrason Tedavisinin Etkinliğinin Karşılaştırılması

Banu Aydeniz,¹ Turan Uslu²

*This study is derived from specialist thesis completed at Fatih Sultan Mehmet Training and Research Hospital in 2010.

¹Department Physical Treatment and Rehabilitation Health Sciences University, Bakırköy Dr. Sadi Konuk Training and Research Hospital, İstanbul, Türkiye

²Private Practice, Kadıköy, İstanbul, Türkiye

Cite this article as:

Aydeniz B, Uslu T. Comparison of the Efficacy of Low-Level Laser Therapy and Ultrasound Treatment in Patients with Shoulder Pain. Bosphorus Med J 2026;13(2):56-62.

Received: 24.04.2026

Revision: 22.05.2026

Accepted: 22.05.2026

Correspondence:

Dr. Banu Aydeniz,
Department Physical Treatment and Rehabilitation Health Sciences University, Bakırköy Dr. Sadi Konuk Training and Research Hospital, İstanbul, Türkiye

Phone:

+90 506 357 31 82

e-mail:

gulmezbanu@hotmail.com

ABSTRACT

Objectives: Shoulder pain is the most common cause of upper extremity pain. This study aimed to comparatively investigate the effects of low-intensity laser and ultrasound therapy on pain and functionality.

Methods: Our study included 60 patients who presented to our outpatient clinic with shoulder pain and limited range of motion. The study was designed as a randomized, prospective, single-blind trial. Patients were randomly assigned to two groups. Both groups received TENS (transcutaneous electrical nerve stimulation), a hot pack, and medical treatment (diclofenac sodium 75 mg once daily). Ultrasound (US) treatment was administered at a dose of 1.5 W/cm² for 5 minutes, and laser therapy was applied at 2000 Hz, 0.6 J/cm², and 905 nm for 6 minutes. The treatment consisted of 10 sessions over 2 weeks. Evaluations were conducted before treatment, immediately after treatment, and at 3 months. During assessments, VAS (visual analog scale) scores during sleep, movement, and rest and active-passive range of motion (ROM) measurements were used. The Constant and UCLA scoring systems were used for shoulder function assessment.

Results: Both groups demonstrated significant improvements in pain, ROM, and functional scores compared with baseline (p<0.05), with effects maintained at 3 months. No significant between-group differences were observed in VAS pain scores or ROM values (p>0.05). However, the US group showed significantly greater improvement in the Constant score compared with the LLLT group (p<0.05).

Conclusion: This study found that both low-level laser therapy and therapeutic ultrasound were effective in reducing pain and improving range of motion in patients with shoulder pain. Therapeutic ultrasound demonstrated superior efficacy compared with laser therapy in restoring shoulder function. Ultrasound therapy can also be considered an effective and reliable treatment method for shoulder pain.

Keywords: Low-level laser therapy; shoulder pain; ultrasound therapy.

ÖZET

Amaç: Omuz ağrısı, üst ekstremitte ağrılarının en yaygın nedenidir. Bu çalışma, son yıllarda kas-iskelet sistemi ağrılarının tedavisinde sıklıkla kullanılan düşük yoğunluklu lazer ve ultrasonun omuz ağrısının tedavisindeki etkinliğini araştırmayı amaçlamıştır.

Yöntem: Çalışmamıza, polikliniğimize omuz ağrısı ve hareket kısıtlılığı şikâyetiyle başvuran 60 hasta dâhil edildi. Çalışma, randomize, prospektif, tek kör bir çalışma olarak tasarlandı. Hastalar randomize olarak iki gruba ayrıldı. Her iki grup da TENS (Transkutanöz Elektriksel Sinir Stimülasyonu), sıcak kompres ve tıbbi tedavi (günde bir kez 75 mg diklofenak sodyum) aldı. Ultrason (US) tedavisi 5 dakika boyunca 1,5 W/cm² dozunda, lazer tedavisi ise 6 dakika boyunca 2000 Hz, 0,6 J/cm² ve 905 nm’de uygulandı. Tedavi, 2 hafta boyunca 10 seanstan oluştu. Değerlendirmeler tedavi öncesinde, hemen sonrasında ve 3 ay sonra yapıldı. Değerlendirmeler sırasında uyku, hareket ve dinlenme hâllerinde VAS (Görsel Analog Skala) ile aktif-pasif hareket açıklığı (ROM) ölçümleri kullanıldı. Omuz fonksiyonunun değerlendirilmesi için Constant ve UCLA skorum sistemleri kullanıldı.

OPEN ACCESS



This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License.

Bulgular: Her iki grup da başlangıç değerlerine kıyasla ağrı, hareket açıklığı (ROM) ve fonksiyonel skorlarda anlamlı iyileşmeler gösterdi ($p<0,05$) ve bu etkiler 3 ay boyunca devam etti. VAS ağrı skorlarında veya ROM değerlerinde gruplar arasında anlamlı bir fark gözlenmedi ($p>0,05$). Bununla birlikte, US grubu, LLLT grubuna kıyasla Constant skorunda anlamlı derecede daha fazla iyileşme gösterdi ($p<0,05$).

Sonuç: Bu çalışma, hem düşük seviyeli lazer tedavisinin hem de terapötik ultrasonun omuz ağrısı olan hastalarda ağrıyı azaltmada ve hareket açıklığını iyileştirmede etkili olduğunu göstermiştir. Terapötik ultrason, omuz fonksiyonunu geri kazandırmada lazer tedavisine göre üstün etkinlik göstermiştir. Ultrason, lazer tedavisine kıyasla omuz ağrısında etkili ve güvenilir bir tedavi yöntemi olarak düşünülebilir.

Anahtar sözcükler: Düşük yoğunluklu lazer tedavisi; omuz ağrısı; ultrason tedavisi.

Shoulder pain is the most common cause of upper extremity pain and ranks third among musculoskeletal disorders, following low back and neck pain.^[1] The prevalence in the general population ranges from 7% to 26%, with lifetime prevalence reported to reach up to 67%. The annual incidence of new cases is estimated to be 0.9–2.5%, peaking in the middle-aged group (45–64 years). Approximately 50% of patients presenting with shoulder pain in primary care settings continue to report symptoms six months later.^[2] Shoulder pain significantly impairs occupational, social, and athletic activities, adversely affecting the quality of daily life and work performance.

The anatomical structure of the shoulder reveals that the glenohumeral joint, while allowing a wide range of motion, is susceptible to injury due to its inherent compromise between mobility and stability. The rotator cuff consists of the supraspinatus, infraspinatus, teres minor, and subscapularis muscles, which together provide dynamic stability to the shoulder joint. When the tendons of these muscles are subjected to repetitive overhead movements and loading, pathologies such as tendinitis, impingement syndrome, and adhesive capsulitis may develop.^[3] Supraspinatus tendinitis is the most frequently reported pathology among the etiologies of shoulder pain, accounting for approximately 65–70% of all shoulder pain cases.^[4]

Treatment options for shoulder pain include medical therapy, physical therapy modalities, exercise, and surgical intervention when necessary. Conservative approaches—including steroid injections, exercise, manual therapy, and physical therapy modalities—have been reported to achieve 40–80% pain reduction and functional improvement.^[5] Among physical therapy modalities, transcutaneous electrical nerve stimulation (TENS), hot/cold packs, therapeutic ultrasound (US), and low-level laser therapy (LLLT) are the most commonly employed.^[6]

Therapeutic ultrasound is a modality that transmits mechanical energy waves at a frequency of 0.5–3 MHz, producing analgesia, increased tissue elasticity, and accelerated healing

through thermal and mechanical tissue effects. Given its ability to reach deeper tissues, it is frequently preferred in chronic conditions such as fibrosis, contracture, and tendinitis.^[7] LLLT, with its analgesic, biostimulant, and anti-inflammatory properties, has gained increasing use in musculoskeletal disorders. Although its mechanism of action is not fully elucidated, it is understood to promote tissue stimulation through cellular and subcellular mechanisms.^[8]

Both modalities are widely used in the treatment of shoulder pathologies; however, the number of randomized controlled trials comparing their efficacy remains limited. The aim of this study was to compare the short- and mid-term effects of LLLT and US—applied in addition to TENS, hot packs, medical treatment, and exercise—on pain, range of motion (ROM), and functionality in patients with shoulder pain.

Methods

This study included a total of 60 patients aged 30–75 years who presented to the Physical Medicine and Rehabilitation outpatient clinic of Fatih Sultan Mehmet Training and Research Hospital with shoulder pain and restricted motion lasting at least three weeks.

The exclusion criteria were malignancy; systemic diseases (pulmonary fibrotic changes, diabetes mellitus, rheumatological conditions); a history of shoulder fracture or reflex sympathetic dystrophy; and advanced cervical spondylosis or restricted cervical motion.

The study was designed as a randomized, prospective, single-blind trial. Patients were randomized into two groups using the sealed-envelope method. Group 1 received therapeutic US ($n=30$) at 1.5 W/cm² for 5 minutes per session. Group 2 received LLLT ($n=30$) at 2000 Hz, 0.6 J/cm², 5 mW, and 905 nm (Endolaser 422, 905 nm) for 6 minutes per session (2 minutes each over the anterior shoulder, posterior shoulder, and greater tubercle). Both groups concurrently received TENS, hot packs, and exercise therapy. The exercise program—consisting of Codman pendulum exercises, passive-active ROM

exercises, posterior capsule stretching, and strengthening exercises—was provided as a home program. Medical treatment (diclofenac sodium, 75 mg tablet, once daily) was recommended for pain control. The total treatment duration was 10 sessions over two weeks. Patients were evaluated at baseline, post-treatment, and at the 3-month follow-up.

Outcome measures included VAS pain scores at rest, during movement, and at night; active and passive ROM measurements in all planes; and specific shoulder examination tests (Hawkins, Neer, supraspinatus, Speed, and Yergason tests). Functional assessment was performed using the Constant shoulder scoring system and the UCLA (University of California–Los Angeles) shoulder rating scale.

The Constant score is widely used to evaluate shoulder function before and after both conservative and surgical interventions and was first described by Constant and Murley in 1986.^[9] The UCLA scale was developed by Ellman in 1986 and is favored for its ease of application.^[10]

Ethical Approval

This study was based on a specialist thesis completed in 2010. At the time the study was initiated, Fatih Sultan Mehmet Training and Research Hospital did not have an institutional ethics committee. Therefore, the study protocol was submitted to and approved by the Fatih Sultan Mehmet Training and Research Hospital Thesis Committee under the petition dated August 26, 2009, and numbered 7653. The study was conducted in accordance with the principles of the Declaration of Helsinki. Written informed consent was obtained from all participants before enrollment.

Statistical Analysis

Statistical analyses were performed using SPSS version 17.0. The chi-square test was used for categorical variables, and

the independent-samples t-test was used for continuous variables. Within- and between-group comparisons of pre- and post-treatment measurements were conducted using repeated-measures two-way analysis of variance and Cochran's Q test. Statistical significance was set at $p \leq 0.05$.

Results

Sixty patients were included in the study; 66.7% ($n=40$) were female, and 33.3% ($n=20$) were male. The mean age was 53.4 years. No statistically significant differences were found between the US and LLLT groups in terms of age, sex, symptom duration, or baseline measurement values ($p > 0.05$) (Table 1). Right-hand dominance was present in 93.3% of patients; however, no significant association was observed between hand dominance and the side of the painful shoulder (Table 2). The groups were statistically comparable at baseline.

Regarding diagnosis distribution, 68.3% ($n=41$) had supraspinatus tendinitis, 8.3% ($n=5$) had biceps tendinitis, 20% ($n=12$) had adhesive capsulitis, and 3.3% ($n=2$) had acromioclavicular joint degeneration. Supraspinatus tendinitis was the most common diagnosis in this study, consistent with the existing literature on shoulder pain etiology.^[3,4] (Table 3).

VAS pain scores (at rest, during movement, and at night) showed statistically significant decreases in both groups compared with baseline following treatment ($p = 0.00 < 0.05$). Improvements observed at the 3-month follow-up also remained significant compared with baseline in both groups. Between-group comparisons revealed no statistically significant differences in VAS rest, movement, or night pain scores between the LLLT and US groups ($p > 0.05$), suggesting comparable analgesic efficacy of both modalities.^[11] (Table 4).

Active and passive ROM measurements—including flexion, abduction, internal rotation, and external rotation—showed significant improvements from baseline in both

Table 1. Comparison of baseline continuous demographic characteristics between groups

Variable	All patients (n=60)		Ultrasound group (n=30)		Laser group (n=30)		p
	Mean	SD	Mean	SD	Mean	SD	
Age (Years)	53.4	11.3	54.2	10.6	52.6	12.1	0.60
Duration (month)	8.6	8.8	8.1	9.3	9.2	8.3	0.64
Weight (kg)	75.0	14.4	72.4	12.9	77.5	15.7	0.17
Height (cm)	162.7	8.9	160.8	8.3	164.6	9.2	0.99

Independent samples t-test

Table 2. Comparison of baseline categorical demographic characteristics between groups

Variable		All patients		Ultrasound group		Laser group		p
		n	%	n	%	n	%	
Gender	Female	40	66.7	23	76.7	17	56.7	0.10
	Male	20	33.3	7	23.3	13	43.3	
Dominant side	Right	56	93.3	29	96.7	27	90	0.10
	Left	4	6.7	1	3.3	3	10	
Affected shoulder	Right	29	48.3	15	50.0	14	46.7	0.80
	Left	31	51.7	15	50.0	16	53.3	
Marital status	Single	5	8.3	1	3.3	4	13.3	0.37
	Married	51	85.0	27	90.0	24	80.0	
	Widowed	4	6.7	2	6.7	2	6.7	
Education level	Illiterate	13	21.7	7	23.3	6	20	0.79
	Primary school	33	55	17	56.7	16	53.3	
	Middle school	4	6.7	2	6.7	2	6.7	
	High school	7	11.7	2	6.7	5	16.7	
	University	3	5	2	6.72	1	3.3	
Employment status	Housewife	34	56.7	19	63.3	15	50	0.48
	Retired	15	25	6	20	9	30	
	Employed	10	16.7	4	13.3	6	20	
	Desk job	1	1.7	1	3.3	0	0	

Chi-square test

Table 3. Distribution of clinical diagnoses between groups

Diagnosis	Ultrasound group n / %	Laser group n / %	Total n / %
Supraspinatus tendinitis	23/76.7%	18/60.0%	41/68.3%
Biceps tendinitis	3/10.0%	2/6.7%	5/8.3%
Adhesive capsulitis	4/13.3%	8/26.7%	12/20.0%
Akromioclavicular joint degeneration	0/0.0%	2/6.7%	2/3.3%
Total	30/100%	30/100%	60/100%

groups ($p < 0.05$). ROM gains achieved after treatment were maintained at the 3-month follow-up. No statistically significant between-group differences were found in active or passive ROM values ($p > 0.05$). Ultrasound's capacity to heat deep tissues is known to increase collagen elasticity and reduce capsular restriction; this biomechanical mechanism is thought to contribute to ROM recovery, particularly in adhesive capsulitis.^[7] (Table 4).

Functional outcomes were assessed using the Constant score and the UCLA shoulder rating scale. UCLA scores showed significant improvement in both groups following treatment; however, the between-group difference was not

statistically significant ($p > 0.05$). In contrast, the Constant score demonstrated statistically significantly greater functional recovery in the US group compared with the LLLT group ($p < 0.05$) (Table 4). This finding indicates that while pain reduction and ROM improvement were comparable between groups, therapeutic US showed a relative advantage in functional recovery. The tissue-heating and fibrolytic properties of ultrasound may play a key role in this effect.^[6]

Discussion

The shoulder is a complex joint that relies on soft-tissue stabilizers—including the rotator cuff muscles, joint capsule,

Table 4. Comparison of VAS, range of motion, and functional scores over time

Outcome	Ultrasound group			Laser group			p
	Pre treatment	Post treatment	3. months	Pre treatment	Post treatment	3. months	
VAS Rest	4.2	2.2	1.2	3.6	2.2	1.6	0.55
VAS Movement	7.6	4.2	3.2	7.1	4.2	2.1	0.31
VAS Sleep	6.3	3.2	2.7	7.9	3.7	2.4	0.16
Abduction (Abd) AROM (°)	125.6	145.0	154.3	127.7	144.6	143.0	0.17
Abd. PROM (°)	132.9	156.3	160.3	132.6	149.7	144.3	0.14
Flexion AROM (°)	145.5	166.1	170.0	150.0	157.7	161.3	0.07
Flexion PROM (°)	153.4	169.5	172.0	153.9	160.8	162.6	0.17
Internal rotation (IR) AROM (°)	61.5	73.2	78.2	64.4	75.2	77.8	0.81
IR PROM (°)	65.3	74.8	79.3	65.2	75.4	77.8	0.87
External rotation (ER) AROM (°)	55.7	67.4	73.0	58.6	65.3	71.2	0.54
ER PROM (°)	61.7	70.7	75.1	60.0	66.6	71.4	0.80
Constant Score	42.4	64.0	71.0	50.5	65.4	67.1	0.01*
UCLA Score	14.7	24.4	27.3	15.6	24.8	25.7	0.23

Two-way ANOVA and Cochran's Q test. AROM: Active range of motion; PROM: Passive range of motion; VAS: Visual analog scale

and ligaments—to achieve its wide range of motion, making it susceptible to injury due to limited bony stabilization. Shoulder pain is the most common cause of upper extremity disorders, with a lifetime prevalence of approximately 70%, leading to significant functional impairment and loss of work capacity.^[12]

Multiple factors contribute to the etiology of shoulder pain. Higher rates are observed in individuals performing frequent overhead activities, such as athletes, swimmers, and weightlifters.^[13] Age is a factor in etiology: glenohumeral instability and rotator cuff tendinitis are more prevalent in individuals under 40 years of age, while rotator cuff tendinitis, adhesive capsulitis, and glenohumeral osteoarthritis are more common after the age of 40.^[14]

The most common causes of shoulder pain include subacromial impingement syndrome, rotator cuff tendinopathy, biceps tendon pathologies, adhesive capsulitis, and acromioclavicular joint disorders.^[15] In this study, 68.3% of patients had supraspinatus tendinitis, 20% had adhesive capsulitis, 8.3% had biceps tendinitis, and 3.3% had acromioclavicular joint degeneration.

The mean age of patients in our study was 53.4 years, with comparable age distributions in both groups, and age-related shoulder pathologies were consistent with the literature. Female patients predominated (66.7%), which is in accordance with studies on the epidemiology of shoulder pain,

supporting the role of hormonal factors and occupational loading in increasing the risk of shoulder pathology in women.^[2] No significant association was observed between hand dominance and the side of the symptomatic shoulder, which is consistent with previous studies reporting inconsistent findings regarding the relationship between shoulder pain and hand dominance, with several demonstrating no significant association between symptom side and the dominant extremity.^[1,16]

Conservative approaches are the first-line treatment for shoulder pain, including steroid injections, exercise, manual therapy, and physical therapy modalities. Given the paucity of studies directly comparing LLLT and therapeutic US in shoulder pain, this study was designed to evaluate and compare the effects of these two modalities on pain, ROM, and functionality. Both modalities produced clinically meaningful improvements; therapeutic US demonstrated statistically significant superiority over LLLT in shoulder functionality ($p=0.01$). Treatment efficacy was maintained at the 3-month follow-up.

A systematic review evaluating the efficacy of LLLT in shoulder pain confirmed that conservative treatment remains the primary approach; however, it emphasized that the level of evidence for the use of passive treatment modalities independently of exercise remains insufficient.^[17] In this context, it is noteworthy that more clinically meaningful im-

provements can be achieved when both modalities are used in conjunction with active exercise. In the present study, exercise therapy was administered concurrently to both groups.

When systematic reviews evaluating the efficacy of LLLT in shoulder pain are examined, it is evident that LLLT applied at appropriate doses provides clinically significant pain relief and that its efficacy increases when combined with physiotherapy.^[17,18] The analgesic mechanism of LLLT is thought to occur through neuronal inhibition, mitochondrial activation, suppression of prostaglandin synthesis, and enhancement of endorphin release.^[8] In light of these mechanisms, the similar degree of pain reduction observed in both treatment groups in our study is consistent with expectations.

A 2024 systematic review and meta-analysis on the efficacy of ultrasound in shoulder pain reported that ultrasound reduces shoulder pain.^[19] One study with controversial findings reported that US therapy, when applied as an adjunct to other physical therapy modalities, provided no significant additional benefit over placebo; however, it was emphasized that, when used alone, it supports tissue healing through thermal and mechanical effects.^[20] We believe that the application of ultrasound within a multimodal regimen alongside TENS and hot packs in our study may have contributed to the superior effect of this combination on functionality. The significant functional superiority observed in US-treated patients at follow-up (third month) compared with those receiving LLLT may be explained by therapeutic ultrasound's capacity to more directly affect the capsular structures of the shoulder joint. The deep tissue heating effect of US softens collagen fibers, increases capsular flexibility, and facilitates synovial fluid production. These mechanisms make important contributions to the restoration of joint function, particularly in adhesive capsulitis and rotator cuff tendinitis.^[21]

A review of the literature revealed that studies directly comparing laser and ultrasound in shoulder pain are limited in number. In a randomized controlled study, LLLT and US therapy were compared in combination with a home exercise program in patients with subacromial impingement syndrome; both modalities yielded effective outcomes regarding pain and functionality, but no statistically significant difference was detected between the groups.^[12] In another study involving patients diagnosed with rotator cuff tendinopathy, similar findings were reported; both the LLLT

and US groups demonstrated marked improvement in pain, function, sleep quality, and emotional status following treatment.^[5]

One of the most important findings of our study is that, while pain reduction and increased range of motion were achieved with both modalities, no significant difference was detected between the groups. This result suggests that the analgesic and kinetic efficacy of LLLT and US in shoulder pain are comparable. Similarly, in a randomized controlled study conducted by Sen et al.^[12] in patients with subacromial impingement syndrome, no significant difference was found between the LLLT and US groups in terms of VAS and ROM values, and both treatments were reported to yield effective outcomes when combined with home exercise.

The mean symptom duration of 8.6 months in our study indicates that the patient population had characteristics of chronic shoulder pain. Based on these findings, both LLLT and therapeutic US can be considered effective interventions for chronic shoulder pain and dysfunction. Current evidence indicates that exercise-based physical therapy approaches have positive effects on pain and function in chronic shoulder pain; combined therapies have been shown to be more effective, while passive modalities alone offer limited benefit.^[22] The results of this study further demonstrate that LLLT or US can be added to the existing treatment regimen in shoulder pain management.

Strengths of this study include its randomized, prospective, single-blind design and the simultaneous use of two validated functional scoring systems. Limitations include a relatively small sample size, the delivery of exercise as a home program rather than a supervised protocol, and the evaluation of treatment efficacy across a heterogeneous group of shoulder pathologies. Future research is recommended with larger sample sizes, active exercise control groups, and investigation of more specific shoulder pathology subgroups.

Conclusion

This study demonstrated that both low-level laser therapy and therapeutic ultrasound are effective in reducing pain and improving the range of motion in patients with shoulder pain. Therapeutic ultrasound showed superior efficacy over laser therapy in restoring shoulder functionality.

These findings indicate that therapeutic ultrasound can be used as a reliable and effective alternative to laser therapy

in the conservative management of shoulder pain. In particular, ultrasound therapy represents a viable option when laser therapy is contraindicated or when cost constraints are a consideration in clinical settings.

Disclosure

Ethics Committee Approval: This study was based on a specialist thesis completed in 2010. At the time the study was initiated, Fatih Sultan Mehmet Training and Research Hospital did not have an institutional ethics committee. Therefore, the study protocol was submitted to and approved by the Fatih Sultan Mehmet Training and Research Hospital Thesis Committee under the petition dated August 26, 2009, and numbered 7653.

Informed Consent: Written informed consent was obtained from all participants included in this study.

Conflict of Interest: None declared.

Financial Disclosure: This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Use of AI for Writing Assistance: The authors declare that no artificial intelligence (AI) or generative ai tools were used in the conception, preparation, writing, analysis, revision, or editing of this manuscript.

Authorship Contributions: Concept – B.A., T.U.; Design – B.A., T.U.; Supervision – B.A., T.U.; Resource – B.A., T.U.; Materials – B.A., T.U.; Data collection and/or processing – B.A., T.U.; Analysis and/or interpretation – B.A., T.U.; Literature review – B.A., T.U.; Writing – B.A., T.U.; Critical review – B.A., T.U.

Peer-review: Externally peer-reviewed.

References

1. Luime JJ, Koes BW, Hendriksen IJ, Burdorf A, Verhagen AP, Miedema HS, et al. Prevalence and incidence of shoulder pain in the general population; a systematic review. *Scand J Rheumatol* 2004;33:73–81.
2. Lucas J, van Doorn P, Hegedus E, Lewis J, van der Windt D. A systematic review of the global prevalence and incidence of shoulder pain. *BMC Musculoskelet Disord* 2022;23:1073.
3. Iordan DA, Leonard S, Matei DV, Sardaru DP, Onu I, Onu A. Understanding scapulohumeral periarthritis: a comprehensive systematic review. *Life (Basel)* 2025;15:186.
4. Spargoli G. Supraspinatus tendon pathomechanics: A current concepts review. *Int J Sports Phys Ther* 2018;13:1083–94.
5. Diyarbakır Ş, Serdaroğlu Beyazal M, Devrimel G, Yıldırım M, Topaloğlu MS. Comparison of the effectiveness of low-level laser therapy and therapeutic ultrasound in patients with rotator cuff tendinopathy. *J Clin Med* 2025;14:4197.
6. Klintberg IH, Cools AM, Holmgren TM, Holzhausen AC, Johansson K, Maenhout AG, et al. Consensus for physiotherapy for shoulder pain. *Int Orthop* 2015;39:715–20.
7. Yildirim MA, Ones K, Celik EC. Comparison of ultrasound therapy of various durations in the treatment of subacromial impingement syndrome. *J Phys Ther Sci* 2013;25:1151–4.
8. El Tayeb MN, Hassan AA, Abuelnaga YG, Eid PM, Tarkhan YM, Fathallah MM, et al. Low level laser therapy versus ultrasonic therapy for the treatment of shoulder impingement syndrome: A systematic review of randomized controlled trials. *QJM: Int J Med* 2021;114(Suppl 1):hcab116.
9. Katolik LI, Romeo AA, Cole BJ, Verma NN, Hayden JK, Bach BR. Normalization of the Constant score. *J Shoulder Elbow Surg* 2005;14:279–85.
10. Constant CR, Murley AH. A clinical method of functional assessment of the shoulder. *Clin Orthop Relat Res* 1987;160–4.
11. Sen EI, Arman S, Tseveendorj N, Yilmaz E, Oral A, Capan N. Low-level laser therapy versus ultrasound therapy combined with home-based exercise in patients with subacromial impingement syndrome: A randomized-controlled trial. *Turk J Phys Med Rehabil* 2023;69:424–33.
12. Sidhar K, Lim HJ, Gutierrez L. A simplified approach to evaluate and manage shoulder Pain. *J Am Board Fam Med* 2024;37:1156–66.
13. Schwarzkopf R, Oron A, Loebenberg M. Shoulder pain: assessment, diagnosis and treatment of common problems. *Harefuah* 2008;147:71–6, 93. [Article in Hebrew]
14. Burbank KM, Stevenson JH, Czarnecki GR, Dorfman J. Chronic shoulder pain: part I. Evaluation and diagnosis. *Am Fam Physician* 2008;77:453–60.
15. Pieters L, Lewis J, Kuppens K, Jochems J, Bruijstens T, Joossens L, et al. An update of systematic reviews examining the effectiveness of conservative physical therapy interventions for subacromial shoulder pain. *J Orthop Sports Phys Ther* 2020;50:131–41.
16. Gill TK, Shanahan EM, Allison D, Alcorn D, Hill CL. Prevalence of abnormalities on shoulder MRI in symptomatic and asymptomatic older adults. *Int J Rheum Dis* 2014;17:863–71.
17. Haslerud S, Magnussen LH, Joensen J, Lopes-Martins RA, Bjordal JM. The efficacy of low-level laser therapy for shoulder tendinopathy: a systematic review and meta-analysis of randomized controlled trials. *Physiother Res Int* 2015;20:108–25.
18. Castaldo M, D'Ossat AA, Gnessi P, Galeoto G. A systematic review on low-level laser therapy in the management of shoulder impingement syndrome. *Appl Sci* 2023;13:3536.
19. Guan H, Wu Y, Wang X, Liu B, Yan T, Abedi-Firouzjah R. Ultrasound therapy for pain reduction in musculoskeletal disorders: a systematic review and meta-analysis. *Ther Adv Chronic Dis* 2024;15:20406223241267217.
20. Kurtaiş Gürsel Y, Ulus Y, Bilgiç A, Dinçer G, van der Heijden GJ. Adding ultrasound in the management of soft tissue disorders of the shoulder: a randomized placebo-controlled trial. *Phys Ther* 2004;84:336–43.
21. Uysal B, Özkuk K, Şahin N, Ökmen BM, Sezer R, Ateş Z. Comparison of the effectiveness of high-intensity laser and ultrasound therapies in adhesive capsulitis: A randomized controlled study. *J Back Musculoskelet Rehabil* 2023;36:227–36.
22. Yang F, Li X, Wang J, Gao Q, Pan M, Duan Z, et al. Efficacy of different analgesic strategies combined with conventional physiotherapy program for treating chronic shoulder pain: a systematic review and network meta-analysis. *J Orthop Surg Res* 2024;19:544.



Effects of Greater Occipital Nerve Block on Headache Frequency and Sleep Disturbance in Medication Overuse Headache

Büyük Oksipital Sinir Bloğunun İlaç Aşırı Kullanım Baş Ağrısında Baş Ağrısı Sıklığı ve Uyku Bozukluğu Üzerine Etkileri

Ahmet Akpınar

ABSTRACT

Objectives: Medication Overuse Headache (MOH) is a secondary headache that develops due to the use of analgesic treatment for more than three months, exceeding 10–15 days per month, depending on the medication used. Patients with MOH are more likely to experience sleep disorders, such as chronic insomnia, compared with those without such disorders, which negatively impacts their quality of life.

Methods: This study was a retrospective, single-center observational study conducted at the Neurology Clinic involving 23 patients diagnosed with MOH and treated with a greater occipital nerve (GON) block. Patients over 18 years of age who met the ICHD-3 criteria for MOH were included in the study. For each patient, the average monthly headache frequency over the three months following the injection, Epworth Sleepiness Scale (ESS) scores, and subjective sleep quality assessments were recorded.

Results: Among the participants, the number of days with pain decreased in 18, increased in 1, and remained the same in 4. A statistically significant difference was found when comparing the number of painful days before and after the procedure. In 7 participants, the ESS score decreased after the procedure, while in 16, it remained the same. A statistically significant difference was found when comparing the ESS scores before and after the procedure.

Conclusion: GON block is an effective, low-cost, accessible, and safe procedure that can be used in the treatment of MOH. Our findings demonstrate a reduction in the number of headache days and daytime sleepiness.

Keywords: GON block; headache; insomnia; medication overuse headache.

ÖZET

Amaç: İlaç Aşırı Kullanım Baş Ağrısı (İAKBA), kullanılan ilaca bağlı olarak ayda 10–15 günü aşacak şekilde, üç aydan uzun süre analjezik tedavi kullanımına bağlı olarak gelişen sekonder bir baş ağrısıdır. İAKBA'lı hastalar, bu tür bozuklukları olmayan kişilere kıyasla kronik uykusuzluk gibi uyku bozukluklarını daha sık yaşamakta olup bu durum yaşam kalitelerini olumsuz yönde etkilemektedir.

Yöntem: Bu çalışma, Nöroloji Kliniği'nde yürütülen, İAKBA tanısı almış ve büyük oksipital sinir (GON) bloğu uygulanmış 23 hastayı kapsayan retrospektif, tek merkezli, gözlemsel bir çalışmadır. İAKBA için ICHD-3 kriterlerini karşılayan 18 yaş üstü hastalar çalışmaya dahil edilmiştir. Her hasta için enjeksiyonu izleyen üç ay boyunca ortalama aylık baş ağrısı sıklığı, Epworth Uykululuk Ölçeği (EUÖ) skorları ve subjektif uyku kalitesi değerlendirmeleri kaydedilmiştir.

Bulgular: Katılımcılar arasında ağrılı gün sayısı 18 hastada azalmış, 1 hastada artmış ve 4 hastada aynı kalmıştır. İşlem öncesi ve sonrası ağrılı gün sayıları karşılaştırıldığında istatistiksel olarak anlamlı bir fark bulunmuştur.

Department of Neurology,
Kulu State Hospital,
Konya, Türkiye

Cite this article as:

Akpınar S. Effects of Greater Occipital Nerve Block on Headache Frequency and Sleep Disturbance in Medication Overuse Headache. Bosphorus Med J 2026;13(2):63-68.

Received: 03.06.2026

Revision: 03.06.2026

Accepted: 31.08.2026

Correspondence:

Dr. Ahmet Akpınar,
Department of Neurology,
Kulu State Hospital,
Konya, Türkiye

Phone:

+90 506 512 69 46

e-mail:

ahmet.akpinar94@gmail.com

OPEN ACCESS



This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License.

Katılımcıların 7'sinde EUÖ skoru işlem sonrası azalırken, 16'sında aynı kalmıştır. İşlem öncesi ve sonrası EUÖ skorları karşılaştırıldığında istatistiksel olarak anlamlı bir fark bulunmuştur.

Sonuç: GON bloğu, İAKBA tedavisinde kullanılabilen etkili, düşük maliyetli, erişilebilir ve güvenli bir işlemdir. Bulgularımız, baş ağrısı gün sayısında ve gündüz uyku kalitesinde azalma olduğunu göstermektedir.

Anahtar sözcükler: Baş ağrısı; GON bloğu; ilaç aşırı kullanım baş ağrısı; uykusuzluk.

Medication Overuse Headache (MOH) is a secondary headache that develops due to the use of analgesic treatment for more than three months, exceeding 10–15 days per month, depending on the medication used.^[1] Affecting approximately 60 million people worldwide, MOH is among the top 20 causes of disability globally.^[2] A study conducted in Türkiye in 2012 determined its prevalence to be 2.1% in the general population and 8.2% among migraine patients.^[3] Patients with MOH are more likely to experience sleep disorders, such as chronic insomnia, compared with those without such disorders, which negatively impacts their quality of life. Early identification and treatment of psychiatric comorbidities and sleep disorders in these patients contribute to improved quality of life and functionality.^[4,5]

Greater occipital nerve (GON) block is a frequently preferred procedure among clinicians for primary and secondary headaches due to its simple technique and high efficacy. It is also used in the treatment of MOH.^[6,7] However, there is no established consensus regarding the injection site, the medication used, or the dosage. Typically, the procedure involves injecting bupivacaine or lidocaine into the nerve on the sensitive side.^[8]

In this study, we aimed to evaluate the effects of GON block application on headache frequency and sleep quality in patients with MOH.

Methods

This study was a retrospective, single-center observational study conducted at the Neurology Clinic of Kulu Regional State Hospital between April 1, 2024, and September 1, 2024, involving 23 patients diagnosed with medication overuse headache (MOH) and treated with a greater occipital nerve (GON) block. The study was conducted in accordance with the Declaration of Helsinki and Good Clinical Practice Guidelines and was approved by the Ethics Committee of Konya City Hospital.

Patients over 18 years of age who met the ICHD-3 criteria for MOH were included in the study. A 0.5% bupivacaine injection was administered to the nerve on the side of the

patient's tenderness, targeting a point 2 cm lateral and 2 cm inferior to the external occipital protuberance. Patients who continued follow-up received up to four GON block injections at one-week intervals. Additionally, patients who could not continue the injections for reasons such as relocating to another city or voluntarily discontinuing treatment were also included in the study, and the effects of the number of injections were analyzed.

For each patient, the average monthly headache frequency over the three months following the injection, Epworth Sleepiness Scale (ESS) scores, and subjective sleep quality assessments were recorded and compared with the average values from the three months prior to the injection. During this period, patients were advised to restrict their analgesic use.

Exclusion Criteria:

1. Patients under 18 years of age
2. Patients with a history of another secondary headache
3. Patients receiving psychiatric treatment
4. Patients who did not adhere to analgesic restriction

Statistical Analyses

To describe the distribution of individuals in terms of demographic and clinical characteristics, numbers (n) and percentages (%) were used. The normality of continuous variables in the study was assessed graphically and using the Shapiro–Wilk test. Descriptive statistics for the variables were presented as median (minimum–maximum) values. Cross-tabulations were created to compare subjective sleep quality assessments among individuals experiencing >50% improvement, and numbers (n), percentages (%), and Fisher's exact test statistics were provided. The Mann–Whitney U test was used to compare age and the number of applications according to the change in benefit (>50%) and ESS score. To investigate whether there were differences in the number of painful days and ESS scores between different measurement times (pre-procedure and post-procedure), the Wilcoxon signed-rank test was used. Among individuals experiencing >50% improvement, correlation analyses

Table 1. Demographic and clinical characteristics of study participants

		All cases (n=23)
Age	Median (Min-Max)	48.87±16.87 47.0 (27-88)
Gender, n (%)		
Female		20 (87)
Male		3 (13)
More than 50% benefit, n (%)		
Yes		14 (60.9)
No		9 (39.1)
Adverse effect, n (%)		
Yes		1 (4.3)
No		22 (95.7)
Subjective Sleep Quality Assessment, n (%)		
No change		18 (78.3)
Improved		5 (21.7)

between the change in ESS score and age and number of applications were conducted, and point-biserial correlation coefficients were provided. IBM SPSS Statistics 21.0 (IBM Corp. Released 2012. IBM SPSS Statistics for Windows, Version 21.0. Armonk, NY: IBM Corp.) and MS Excel 2007 were used for statistical analyses and calculations. A statistical significance level of $p < 0.05$ was accepted.

Results

The average age of the participants in the study was 48.87 ± 16.87 years. Of the participants, 87.0% ($n=20$) were women. Among the participants, 60.9% ($n=14$) reported experiencing >50% improvement. One participant (4.3%) experienced an adverse effect, with vasovagal syncope observed. According to the subjective sleep quality assessment, there was no change in 78.3% of the participants ($n=18$) (Table 1).

Table 3. Relationship between more than 50% benefit and number of applications

	More Than 50% Benefit	
	r_{pb}	p
Number of Applications	0.558	0.006

rpb: Point Biserial Correlation Coefficient

Among the participants, the number of days with pain decreased in 18, increased in 1, and remained the same in 4. A statistically significant difference was found when comparing the number of painful days before and after the procedure ($z=3.787$, $p < 0.001$). Additionally, the median number of painful days was 13 before the procedure and 5 after the procedure (Table 2).

According to the point-biserial correlation analysis, there was a positive and statistically significant relationship between experiencing >50% improvement and the number of applications ($p < 0.05$). As the number of applications increased, the likelihood of experiencing >50% improvement also increased (Table 3).

In 7 participants, the ESS score decreased after the procedure, while in 16, it remained the same. A statistically significant difference was found when comparing the ESS scores before and after the procedure ($z=2.375$, $p=0.018$). Additionally, the median ESS score was 6.0 before the procedure and 5 after the procedure (Table 5).

Among individuals who benefited by more than 50%, sleep quality improved in 28.6% ($n=4$), while among those who did not benefit, sleep quality improved in 11.1% ($n=1$). No statistically significant difference was found in the subjective sleep quality assessment according to whether participants experienced >50% improvement ($p=0.327$). In 7 participants, the ESS score decreased after the procedure, while in 16, it remained the same. A statistically significant difference was found when comparing the ESS scores before

Table 2. Comparison of painful days before and after the procedure

Painful days		n	Pre-Procedure Median (Min-Max)	Post-Procedure Median (Min-Max)	z	p
Pre-Procedure - Post-Procedure Change	Increase	1				
	Decrease	18	13 (10-20)	5 (1-20)	$z=3.787$	<0.001
	No change	4				

z: Wilcoxon Signed Rank Test

Table 4. The status of more than 50% benefit based on patients' age

More Than 50% Benefit				
	Yes (n=14)	No (n=9)	Test statistic	
	Median (Min-Max)	Median (Min-Max)	z	p
Age	53 (27-88)	36 (29-75)	z=1.609	0.109
z: Mann Whitney U Test				

and after the procedure ($z=2.375$, $p=0.018$). Additionally, the median ESS score was 6 before the procedure and 5 after the procedure (Table 6).

No statistically significant difference was found in the age or number of applications of the participants according to the change in their ESS scores ($p>0.05$) (Table 7).

Discussion

In this observational study, the effects of greater occipital nerve (GON) block on the number of painful days and sleep quality in patients diagnosed with medication overuse

Table 7. Comparison of age and number of applications based on epworth sleepiness scale

Epworth Sleepiness Scale				
	No Change (n=16)	Decrease (n=7)	Test statistic	
	Median (Min-Max)	Median (Min-Max)	z	p
Age	48.0 (27-88)	37.0 (34-58)	z=0.937	0.376
Number of Applications	2.5 (1-4)	4.0 (1-4)	z=0.932	0.413
z: Mann Whitney U Test				

headache (MOH) were evaluated, and whether increasing the frequency of application improves its effectiveness was investigated.

After the procedure, a statistically significant and notable reduction was observed in the number of painful days. In a study by Karadaş et al.^[9], GON block application using lidocaine three times at weekly intervals was compared with a single application, and both methods were found to be effective. In another study by Arab et al.^[10], GON block was

Table 5. Comparison of epworth sleepiness scale before and after the procedure

Epworth Sleepiness Scale	n	Pre-procedure median (Min-Max)	Post-procedure median (Min-Max)	z	p
Pre-Procedure - Post-Procedure Change	Increase	0			
	Decrease	7	6 (0-12)	5 (0-12)	z=2.375
	No change	16			0.018

z: Wilcoxon Signed Rank Test

Table 6. Comparison of subjective sleep quality assessment according to the status of more than 50% benefit and comparison of epworth sleepiness scale before and after the procedure

More Than %50 Benefit						
		Yes (n=14) n (%)	No (n=9) n (%)		p*	
Subjective Sleep Quality Assessment						
No Change		10 (71.4)	8 (88.9)		0.327	
Improve		4 (28.6)	1 (11.1)			
Epworth Sleepiness Scale		n	Pre-procedure median (Min-Max)	Post-procedure median (Min-Max)	z	p
Pre-Procedure	Increase	0				
Post-Procedure	Decrease	7	6 (0-12)	5 (0-12)	z=2.375	0.018
Change	No change	16				

Fisher Exact test results are provided. z: Wilcoxon Signed Rank Test

also found to be effective in addition to analgesic restriction. In the study by Ünsal et al.^[7], the benefit of GON block application was also demonstrated. Our findings are consistent with the literature in this regard.

There is no standardized procedure regarding the number and frequency of GON blocks. In the study by Karadaş et al.^[9], patients who received a single injection were compared with those who received injections three times at weekly intervals, and repeated injections were shown to be superior. Another study evaluating the effectiveness of repeated GON blocks in cervicogenic headaches also showed the superiority of repeated injections over a single application.^[11] In our study, similar to the literature, it was observed that the number of painful days decreased more with repeated injections. Although a standardized procedure cannot be established based on these findings, repeated applications may be more effective.

In our study, no relationship was found between benefiting from the application and patient age. In a study by Allen et al.^[12], patients were divided into groups based on age (under and over 65 years), and both groups were shown to benefit equally. In another study by Ökmen et al.^[13], no difference in effectiveness was observed between different age groups. These results indicate that our findings are consistent with the literature.

The ESS was used to assess daytime sleepiness in our study, and subjective evaluations of sleep quality were obtained from the patients themselves. A statistically significant decrease in ESS scores was observed after the procedure, but no significant difference was found in subjective sleep quality. Additionally, in patients with a >50% reduction in pain frequency after the injection, a greater decrease in ESS scores was observed, but no changes were observed with an increasing number of injections. In a study by Sağmacı et al.^[14], the sleep quality of the included patients was shown to improve, and psychiatric symptoms regressed. Similar positive effects on sleep quality have been demonstrated in other studies in the literature conducted in patients with chronic migraine.^[15,16] These results suggest that GON block may improve sleep quality and reduce insomnia symptoms.

Limitations

The limitations of this study include its retrospective nature, the subjective assessment of sleep quality, its single-center design, and the small number of patients. The strengths of

the study include directly assessing sleep quality in patients with MOH and evaluating sleep-related issues from the patients' own perspective.

Conclusion

In conclusion, GON block is an effective, low-cost, accessible, and safe procedure that can be used in the treatment of MOH. Our findings demonstrate a reduction in the number of headache days and daytime sleepiness. Although less robust, the findings also suggest an improvement in patients' subjective sleep quality. Prospective, randomized controlled trials with larger sample sizes are needed to expand the available evidence.

Disclosure

Ethics Committee Approval: This study was approved by The Ethics Committee of Konya City Hospital (2025/09 -08.09.2025)

Informed Consent: Written informed consents were obtained from patients who participated in this study.

Conflict of Interest: The authors declare that there is no conflict of interest.

Use of AI for Writing Assistance: The authors declared that artificial intelligence was not used in the study.

Financial Disclosure: The authors declared that this study received no financial support.

Peer-review: Externally peer-reviewed.

References

- Wakerley BR. Medication-overuse headache. *Pract Neurol* 2019;19:399–403.
- Linde M, Gustavsson A, Stovner LJ, Steiner TJ, Barré J, Katsarava Z, et al. The cost of headache disorders in Europe: the Eurolight project. *Eur J Neurol* 2012;19:703–11.
- Ertas M, Baykan B, Orhan EK, Zarifoglu M, Karli N, Saip S, et al. One-year prevalence and the impact of migraine and tension-type headache in Turkey: a nationwide home-based study in adults. *J Headache Pain* 2012;13:147–57.
- Kniazeva IA, Gilev DV, Lebedeva ER. Sleep disorders in patients with medication-overuse headache. *Zh Nevrol Psikhiatr Im S S Korsakova* 2024;124:93–8. [Article in Russian]
- Pompili M, Innamorati M, Lamis DA, Serafini G, Ricci F, Migliorati M, et al. Depression and insomnia are independently associated with satisfaction and enjoyment of life in medication-overuse headache patients. *Int J Psychiatry Med* 2016;51:442–55.
- Chowdhury D, Datta D, Mundra A. Role of greater occipital nerve block in headache disorders: a narrative review. *Neurol India* 2021;69:S228–56.
- Mirac UA, Tugrul A. The effect of anesthetic blockade of greater occipital nerve during the withdrawal period of the medication overuse headache treatment. *Ideggyogy Sz* 2024;77:103–10.

12. Inan LE, Inan N, Unal-Artık HA, Atac C, Babaoglu G. Greater occipital nerve block in migraine prophylaxis: Narrative review. *Cephalalgia* 2019;39:908–20.
13. Karadaş Ö, Özön AÖ, Özçelik F, Özge A. Greater occipital nerve block in the treatment of triptan-overuse headache: A randomized comparative study. *Acta Neurol Scand* 2017;135:426–33.
14. Arab A, Khoshbin M, Karimi E, Saberian G, Saadatnia M, Khorvash F. Effects of greater occipital nerve block with local anesthetic and triamcinolone for treatment of medication overuse headache: an open-label, parallel, randomized, controlled clinical trial. *Neurol Sci* 2022;43:549–57.
15. Naja ZM, El-Rajab M, Al-Tannir MA, Ziade FM, Tawfik OM. Repetitive occipital nerve blockade for cervicogenic headache: expanded case report of 47 adults. *Pain Pract* 2006;6:278–84.
16. Allen SM, Mookadam F, Cha SS, Freeman JA, Starling AJ, Mookadam M. Greater occipital nerve block for acute treatment of migraine headache: a large retrospective cohort study. *J Am Board Fam Med* 2018;31:211–8.
17. Okmen K, Dagistan Y, Dagistan E, Kaplan N, Cancan E. Efficacy of the greater occipital nerve block in recurrent migraine type headaches. *Neurol Neurochir Pol* 2016;50:151–4.
18. Saçmacı H, Aktürk T, Tanık N, Inan LE. Clinical success of greater occipital nerve blockade in improving sleep quality of chronic migraineurs: a prospective observational study. *Sleep Breath* 2021;25:1003–10.
19. Metin KM, Dilek SS, Karaduman Y, Serçe A, Takmaz SA, Inan LE. The effects of greater occipital nerve blockage with lidocaine on sleep characteristics in chronic migraine patients. *Clin Neurol Neurosurg* 2023;231:107826.
20. Şahin O, Güneş M, Şahin EHK, Dönmez R. The effect of greater occipital nerve block on amplification of somatic symptoms and anxiety and depression levels in chronic migraine. *Acta Neurol Belg* 2024;124:1641–6.



An Early-Stage Neuroborreliosis Case with Atypical Findings

Atipik Bulgularıyla Erken Dönem Nöroborrelioz Olgusu

✉ Sema Yapıcı, ✉ Elif Söylemez, ✉ Mehmet Fatih Özdağ, ✉ Sibel Karşıdağ

ABSTRACT

Lyme disease is a zoonotic infection caused by the spirochete *Borrelia burgdorferi*, which can affect multiple organs, including the skin, joints, heart, and central nervous system. Neurological involvement that typically appears within the first few months after a tick bite is referred to as neuroborreliosis. In this paper, we present a case of neuroborreliosis with concurrent involvement of the trigeminal nerve, peripheral facial nerve, and partial cauda equina. A 32-year-old male patient presented with complaints of facial numbness, difficulty walking, and loss of sensation in the perianal region. Brain and spinal magnetic resonance imaging (MRI) revealed no pathological findings. Cerebrospinal fluid (CSF) analysis showed elevated protein levels, and *Borrelia burgdorferi* IgM antibodies were positive. After the patient was treated with intravenous ceftriaxone, significant improvement in clinical symptoms was observed. This case is important because it demonstrates that neuroborreliosis can present with rare neurological findings, such as cranial neuropathy and cauda equina syndrome.

Keywords: Cauda equina syndrome; cranial neuropathy; Lyme disease; neuroborreliosis.

ÖZET

Lyme hastalığı, *Borrelia burgdorferi* isimli spiroket tarafından oluşturulan ve deri, eklem, kalp ile sinir sistemi başta olmak üzere birçok organı etkileyebilen zoonotik bir enfeksiyondur. Kene ısırığından sonra genellikle ilk birkaç ay içinde ortaya çıkan sinir sistemi tutulumu nöroborrelioz olarak adlandırılır. Bu yazıda, trigeminal sinir, periferik fasiyal sinir ve parsiyel kauda equina tutulumunun birlikte görüldüğü bir nöroborrelioz olgusu sunulmuştur. Otuz iki yaşında erkek hasta, yüzde uyuşma, yürüme güçlüğü ve perianal bölgede his kaybı şikâyetleri ile başvurmuştur. Beyin ve spinal manyetik rezonans görüntüleme incelemelerinde patolojik bulgu saptanmamıştır. Beyin omurilik sıvısı incelemesinde protein yüksekliği saptanmış, *Borrelia burgdorferi* IgM antikorları pozitif bulunmuştur. Hastaya uygulanan intravenöz seftriksone tedavisi sonrasında klinik bulgulara belirgin düzelme izlenmiştir. Bu olgu, nöroborreliozun kraniyal nöropati ve kauda equina sendromu gibi nadir nörolojik bulgular ile prezente olabileceğini göstermesi açısından önem taşımaktadır.

Anahtar sözcükler: Kauda equina sendromu; kraniyal nöropati; Lyme hastalığı; nöroborrelioz.

Department of Neurology,
Sultan Abdülhamid Han
Training and Research
Hospital, İstanbul, Türkiye

Cite this article as:

Yapıcı S, Söylemez E, Özdağ MF, Karşıdağ S. An Early-Stage Neuroborreliosis Case with Atypical Findings. Bosphorus Med J 2026;13(2):69-73.

Received: 25.03.2026

Revision: 17.08.2026

Accepted: 08.09.2026

Correspondence:

Dr. Sema Yapıcı.

Department of Neurology,
Sultan Abdülhamid Han
Training and Research
Hospital, İstanbul, Türkiye

Phone:

+90 538 714 87 47

e-mail:

semayapici1@gmail.com

OPEN ACCESS



This work is licensed under a
Creative Commons Attribution-
NonCommercial 4.0 International
License.

Lyme disease is a multisystemic zoonotic infection transmitted by ticks and caused by the spirochete *Borrelia burgdorferi* sensu lato complex. The primary vectors are *Ixodes* species ticks, and the disease is mainly characterized by skin, joint, cardiac, and nervous system involvement.^[1] The earliest and most typical clinical manifestation is erythema migrans; in later

stages, the disease may lead to neurological, cardiac, and rheumatological complications.^[1]

Nervous system involvement in Lyme disease is termed neuroborreliosis and occurs in approximately 3–12% of cases.^[2] Lyme neuroborreliosis encompasses a broad clinical spectrum that can affect both the central and peripheral nervous systems. The most frequent neurological man-

ifestations include meningoradiculitis, cranial neuropathies, and lymphocytic meningitis.^[3,4]

In this paper, we present a rare case of a patient presenting with facial numbness, facial paralysis, and gait difficulty who was diagnosed with Lyme neuroborreliosis after a thorough evaluation, demonstrating concomitant involvement of cranial nerves V and VII along with partial cauda equina involvement.

Case Report

A 32-year-old male patient presented to the emergency department with left-sided facial numbness, nausea, difficulty walking, perianal numbness, and fecal incontinence. No history of a tick bite could be elicited, although he had a history of recent travel to Georgia. His medical history was unremarkable except for residual visual loss in the right eye.

He had no history of smoking, alcohol, or substance use. On neurological examination, cranial nerve assessment was intact except for residual limitation of right eye abduction (impaired lateral gaze), sensory loss on the left half of the face, and left-sided peripheral facial paralysis. The patient reported no pain. Muscle strength was 4/5 in the proximal lower extremities bilaterally; deep tendon reflexes revealed brisk patellar reflexes bilaterally, normoactive Achilles reflexes, and mute plantar responses. Saddle hypoesthesia starting from the L3 level and encompassing the sacral dermatomes was present. His gait was spastic-ataxic and required support. Routine biochemistry and complete blood count parameters were within normal limits. Serum VDRL and HIV tests were negative. Contrast-enhanced brain, cervical, thoracic, and lumbar spinal magnetic resonance imaging (MRI) studies were normal (Fig.1-4). On electrophysiological eval-




Figure 1. a, b, c. Contrast-enhanced axial T1-weighted images show no evidence of pathological signal intensity involving the brainstem structures. Furthermore, axial FLAIR sequences at the level of the lateral ventricles demonstrate no pathological signal changes within the brain parenchyma.




Figure 2. Contrast-enhanced axial T1-weighted images show no evidence of pathological signal intensity involving the brain-stem structures. Furthermore, axial FLAIR sequences at the level of the lateral ventricles demonstrate no pathological signal changes within the brain parenchyma.

uation, median and tibial somatosensory evoked potentials (SEPs) were within normal limits. Cerebrospinal fluid (CSF) analysis demonstrated elevated protein levels (1.27 g/L) and normal glucose levels (75.5 mg/dL, with a simultaneous blood glucose level of 116 mg/dL); no cells were detected. CSF meningitis panel, CSF culture, tuberculosis polymerase chain reaction (PCR), fungal culture, herpes PCR, *Brucella* IgM and IgG (ELISA), and cytomegalovirus PCR were all negative. CSF oligoclonal band Type 1 was negative. CSF *B. burgdorferi* IgG (Western blot) was negative, whereas CSF *B. burgdorferi* IgM was positive (36.8 Positive U/mL).

During clinical follow-up, his neurological condition progressed: proximal muscle strength in the right lower extremity declined to 3/5, sensory deficits became more prominent, and fecal incontinence developed. Intravenous ceftriaxone,




Figure 3. Contrast-enhanced axial T1-weighted images show no evidence of pathological signal intensity involving the brain-stem structures. Furthermore, axial FLAIR sequences at the level of the lateral ventricles demonstrate no pathological signal changes within the brain parenchyma.



Figure 4. Sagittal T2-weighted lumbar MRI sections demonstrate no pathological signal intensity changes.

2 g once daily, was administered for 3 weeks. The patient's clinical condition gradually improved, and his neurological examination was completely normal upon discharge.

Discussion

In this patient presenting with acute conus-cauda involvement and isolated cranial neuropathy symptoms, a diagnosis of neuroborreliosis was established following comprehensive evaluations. Lyme disease is a multisystemic and multistage infectious disease.^[1]

The clinical features of the disease vary depending on the stage of infection. While cutaneous involvement is most commonly observed in the early stage, systemic organ involvement emerges in later stages. The early objective skin lesion is erythema migrans, whereas chronic cutaneous changes may develop in the late stages.^[1]

Neurological involvement occurs in approximately 3–12% of patients.^[2] Lyme neuroborreliosis can affect both the central and peripheral nervous systems.^[3] In the vast majority of early-stage European neuroborreliosis cases, the clinical presentation manifests as painful meningoradiculitis (Bannwarth syndrome), characterized by severe radicular pain, cranial neuropathy, and CSF pleocytosis.^[4]

Cranial nerve involvement is one of the most common manifestations of Lyme neuroborreliosis, with peripheral facial palsy being particularly frequently reported.^[3,5] Less commonly, the trigeminal nerve and other cranial nerves may also be affected.^[3] Furthermore, lymphocytic meningitis is a common finding in neuroborreliosis, whereas encephalitis or myelitis is observed less frequently.^[3,5]

Cerebral vasculitis is a rare but important complication of Lyme disease and should be considered in the differential diagnosis, especially in young patients presenting with unexplained stroke.^[6] Polyneuropathy, a late-stage manifestation, typically presents as predominantly sensory axonal neuropathy.^[7]

Our patient presented with peripheral facial palsy, trigeminal nerve involvement, and partial cauda equina syndrome. Although cranial nerve involvement is common in neuroborreliosis, the development of a cauda equina-like syndrome concomitant with trigeminal and facial nerve involvement is a clinical combination rarely reported in the literature.^[8]

Inquiring about a history of tick bite is essential for diagnosis; however, only a minority of patients recall prior tick exposure.^[9]

CSF analysis plays a critical role in the diagnostic workup. Protein elevation and inflammatory changes in the CSF can

be observed in cases of neuroborreliosis.^[3] However, culturing the pathogen from CSF is extremely challenging. The sensitivity of PCR testing in CSF is also limited, yielding positive results in only a small subset of cases.^[10]

The current standard approach for the serological diagnosis of Lyme borreliosis is a two-tiered testing algorithm. The first step involves screening with an enzyme immunoassay or immunofluorescence antibody test; positive or equivocal results must be confirmed using immunoblotting.^[10,11] The most specific diagnostic method is the demonstration of intrathecal antibody production specific to *Borrelia burgdorferi* in CSF.^[12]

In our case, although CSF pleocytosis was not detected, elevated protein levels and *B. burgdorferi* IgM positivity were demonstrated by Western blot. While lymphocytic pleocytosis in CSF is observed in the majority of neuroborreliosis cases, cellular expansion may not be detected in early-stage or rare cases. The acute onset of symptoms and the marked clinical response to antibiotic therapy suggested that the detected serological finding was consistent with an active infection.

In the treatment of neuroborreliosis, a 14-day course of antibiotic therapy is considered sufficient for early-stage and isolated neurological involvement, whereas the duration may be extended to 21 days in late-stage or severe central nervous system involvement. First-line treatment frequently involves intravenous ceftriaxone; however, oral doxycycline is known to yield clinical efficacy equivalent to that of intravenous regimens.^[13,14]

Antibiotic therapy leads to complete clinical recovery in the majority of patients; nevertheless, post-treatment Lyme disease syndrome (PTLDS) or persistent symptoms—characterized by ongoing fatigue, widespread musculoskeletal pain, and cognitive dysfunction persisting for months after treatment—may be observed in a subset of patients.^[15]

Conclusion

In conclusion, this case demonstrates that neuroborreliosis can present with atypical neurological features, such as cranial nerve involvement combined with partial cauda equina syndrome. In the presence of clinical suspicion, early serological evaluation and appropriate antibiotic treatment are paramount to the prognosis of the disease.

Disclosure

Informed Consent: Written informed consents were obtained from patients who participated in this study.

Conflict of Interest: The authors declare that there is no conflict of interest.

Use of AI for Writing Assistance: AI-assisted technologies were used for text editing and manuscript checking. During revision, OpenAI Codex was also used to assist with reference verification, revision of the corresponding text, and preparation of this disclosure. The authors remain responsible for the accuracy and integrity of the manuscript.

Financial Disclosure: The authors declared that this study received no financial support.

Authorship Contributions: Concept – M.F.Ö.; Design – E.S., S.Y.; Supervision – S.K., M.F.Ö.; Materials – S.Y.; Data Collection and/or Processing – S.Y.; Analysis and/or Interpretation – S.Y., E.S.; Literature Search – S.Y., E.S.; Writing – S.Y.; Critical Reviews – S.K., M.F.Ö.

Peer-review: Externally peer-reviewed.

References

1. Stanek G, Wormser GP, Gray J, Strle F. Lyme borreliosis. *Lancet* 2012;379:461–73.
2. Kortela E, Kanerva MJ, Puustinen J, Hurme S, Airas L, Lauhio A, et al. Oral doxycycline compared to intravenous ceftriaxone in the treatment of lyme neuroborreliosis: a multicenter, equivalence, randomized, open-label trial. *Clin Infect Dis* 2021;72:1323–31.
3. Halperin JJ. Neuroborreliosis. *Neurol Clin* 2018;36:821–30.
4. Ogrinc K, Lusa L, Lotrič-Furlan S, Bogovič P, Stupica D, Cerar T, et al. Course and outcome of early european lyme neuroborreliosis (Bannwarth Syndrome): clinical and laboratory findings. *Clin Infect Dis* 2016;63:346–53.
5. Knudtzen FC, Andersen NS, Jensen TG, Skarphéðinsson S. Characteristics and clinical outcome of lyme neuroborreliosis in a high endemic area, 1995-2014: a retrospective cohort study in Denmark. *Clin Infect Dis* 2017;65:1489–95.
6. Back T, Grünig S, Winter Y, Bodechtel U, Guthke K, Khatri D, et al. Neuroborreliosis-associated cerebral vasculitis: long-term outcome and health-related quality of life. *J Neurol* 2013;260:1569–75.
7. Kindstrand E, Nilsson BY, Hovmark A, Pirskanen R, Asbrink E. Peripheral neuropathy in acrodermatitis chronica atrophicans - a late Borrelia manifestation. *Acta Neurol Scand* 1997;95:338–45.
8. Bigi S, Aebi C, Nauer C, Bigler S, Steinlin M. Acute transverse myelitis in Lyme neuroborreliosis. *Infection* 2010;38:413–6.
9. Henningsson AJ, Malmvall BE, Ernerudh J, Matussek A, Forsberg P. Neuroborreliosis-an epidemiological, clinical and healthcare cost study from an endemic area in the south-east of Sweden. *Clin Microbiol Infect* 2010;16:1245–51.
10. Branda JA, Steere AC. Laboratory diagnosis of lyme borreliosis. *Clin Microbiol Rev* 2021;34:e00018–19.
11. Mead P, Petersen J, Hinckley A. Updated CDC recommendation for serologic diagnosis of lyme disease. *MMWR Morb Mortal Wkly Rep* 2019;68:703.
12. Lantos PM, Rumbaugh J, Bockenstedt LK, Falck-Ytter YT, Agüero-Rosenfeld ME, Auwaerter PG, et al. Clinical Practice Guidelines by the Infectious Diseases Society of America, American Academy of Neurology, and American College of Rheumatology: 2020 Guidelines for the Prevention, Diagnosis, and Treatment of Lyme Disease. *Neurology* 2021;96:262–73.
13. Rauer S, Kastenbauer S, Hofmann H, Fingerle V, Huppertz HI, Hunfeld KP, et al. Guidelines for diagnosis and treatment in neurology - Lyme neuroborreliosis. *Ger Med Sci* 2020;18:Doc03.
14. Ljøstad U, Skogvoll E, Eikeland R, Midgard R, Skarpaas T, Berg A, et al. Oral doxycycline versus intravenous ceftriaxone for European Lyme neuroborreliosis: a multicentre, non-inferiority, double-blind, randomised trial. *Lancet Neurol* 2008;7:690–5.
15. Rebman AW, Aucott JN. Post-treatment lyme disease as a model for persistent symptoms in lyme disease. *Front Med (Lausanne)* 2020;7:57.

